

148-1276

**AGRICULTURAL**

**ALFALFA** 8 ster beetes... 5 to 15 per acre...  
**APPLES** 10 ster beetes... 10 to 20 per acre...  
**APRICOTS** 10 ster beetes... 10 to 20 per acre...  
**ASPARAGUS** 10 ster beetes... 10 to 20 per acre...  
**BEANS** 10 ster beetes... 10 to 20 per acre...  
**BLACKBERRIES** 10 ster beetes... 10 to 20 per acre...  
**BLUEBERRIES** 10 ster beetes... 10 to 20 per acre...  
**BOYSENBERRIES** 10 ster beetes... 10 to 20 per acre...  
**BROCCOLI** 10 ster beetes... 10 to 20 per acre...  
**BRUSSELS SPROUTS** 10 ster beetes... 10 to 20 per acre...  
**CABBAGE** 10 ster beetes... 10 to 20 per acre...  
**CARROTS** 10 ster beetes... 10 to 20 per acre...  
**CAULIFLOWER** 10 ster beetes... 10 to 20 per acre...  
**CHERRIES** 10 ster beetes... 10 to 20 per acre...  
**CHINESE CABBAGE** 10 ster beetes... 10 to 20 per acre...  
**CUCUMBER** 10 ster beetes... 10 to 20 per acre...  
**COLLARDS** 10 ster beetes... 10 to 20 per acre...  
**CORN** 10 ster beetes... 10 to 20 per acre...  
**CORN MEAL** 10 ster beetes... 10 to 20 per acre...  
**CORNFLOUR** 10 ster beetes... 10 to 20 per acre...  
**CORNFLOUR** 10 ster beetes... 10 to 20 per acre...

**PARSLEY** 10 ster beetes... 10 to 20 per acre...  
**PARSNIPS** 10 ster beetes... 10 to 20 per acre...  
**PASTURE AND RANGELAND** 10 ster beetes... 10 to 20 per acre...  
**PEACHES** 10 ster beetes... 10 to 20 per acre...  
**PEANUTS** 10 ster beetes... 10 to 20 per acre...  
**PEARS** 10 ster beetes... 10 to 20 per acre...  
**PEPPERS** 10 ster beetes... 10 to 20 per acre...  
**PLUMS** 10 ster beetes... 10 to 20 per acre...  
**POTATOES** 10 ster beetes... 10 to 20 per acre...  
**PRUNES** 10 ster beetes... 10 to 20 per acre...  
**PUMPKINS** 10 ster beetes... 10 to 20 per acre...  
**RADISHES** 10 ster beetes... 10 to 20 per acre...  
**RASPBERRIES** 10 ster beetes... 10 to 20 per acre...  
**RICE** 10 ster beetes... 10 to 20 per acre...  
**RUTABAGA** 10 ster beetes... 10 to 20 per acre...  
**SALAD** 10 ster beetes... 10 to 20 per acre...  
**SORGHUM** 10 ster beetes... 10 to 20 per acre...  
**SWEETANS** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...  
**TANSENG** 10 ster beetes... 10 to 20 per acre...

BEST DOCUMENT AVAILABLE

148-1274

(2)

usually 30 to 35 feet is indicated. For application with ground equipment, use 100 to 150 lb per acre with a spraying boom 25 to 30 feet wide. For hand application, use 50 to 75 lb per acre. Use the higher application rates for rough ground. Grasshoppers—10 to 15 lb per acre. Use the higher application rate for rough ground. Beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Strawberry rootworm—10 to 15 lb per acre. Refer to separate recommendations for more details. Strawberry rootworm—10 to 15 lb per acre. Refer to separate recommendations for more details.

**CUCUMBERS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Cucumber beetle—10 to 15 lb per acre. Refer to separate recommendations for more details. Yellow fever fever—10 to 15 lb per acre. Refer to separate recommendations for more details. Weevils—10 to 15 lb per acre. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per acre. Refer to separate recommendations for more details.

**CUCUMBERS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Cucumber beetle—10 to 15 lb per acre. Refer to separate recommendations for more details. Yellow fever fever—10 to 15 lb per acre. Refer to separate recommendations for more details. Weevils—10 to 15 lb per acre. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per acre. Refer to separate recommendations for more details.

**CUCUMBERS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Cucumber beetle—10 to 15 lb per acre. Refer to separate recommendations for more details. Yellow fever fever—10 to 15 lb per acre. Refer to separate recommendations for more details. Weevils—10 to 15 lb per acre. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per acre. Refer to separate recommendations for more details.

**CUCUMBERS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Cucumber beetle—10 to 15 lb per acre. Refer to separate recommendations for more details. Yellow fever fever—10 to 15 lb per acre. Refer to separate recommendations for more details. Weevils—10 to 15 lb per acre. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per acre. Refer to separate recommendations for more details.

**TOMATOES**—14 day pre-harvest interval—Colorado potato beetle—10 to 15 lb per acre. Refer to separate recommendations for more details. Flea beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per acre. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per acre. Refer to separate recommendations for more details.

**POULTRY (Chickens, Ducks, Game Birds, Geese, Partridges, Pheasants, Turkeys)**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per acre. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per acre. Refer to separate recommendations for more details.

**POULTRY HOUSES**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per acre. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per acre. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per acre. Refer to separate recommendations for more details.

**PEST CONTROL**  
**DOMESTIC DWELLINGS**—Earwigs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Fleas—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Fleas—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Fleas—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Fleas—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Fleas—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Fleas—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Fleas—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

**ALL RATES SHOULD BE APPLIED AS FOLIAGE APPLICATIONS UNLESS OTHERWISE NOTED**

**ORNAMENTALS**  
**LANNS AND TURFS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

**HERBACEOUS PLANTS AND BULBS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

**WOODY SHRUBS, TREES AND VINES**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

**NONAGRICULTURAL LANDS, PASTURE RANGELAND**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

**ROCKED AREAS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

**URBAN AND RURAL OUTDOOR AREAS (PUBLIC HEALTH)**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

**DOGHOUSES, KENNELS, SLEEPING QUARTERS**—14 day pre-harvest interval—Flea beetles—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Leafhoppers—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details. Nymphs—10 to 15 lb per 500 square feet. Refer to separate recommendations for more details.

# PRO SEAMAN<sup>®</sup> 100% DUST

ACCEP  
JAN-12

148-12

**BEST DOCUMENT AVAILABLE**

ACTIVE INGREDIENTS:	By Weight
1-naphthyl-N-methylcarbamate	10
INERT INGREDIENTS	90
TOTAL	100

ST. VIN. 100-1000-1 Made in U.S.A. by Union Carbide Corporation. For the active ingredient: 1-naphthyl-N-methylcarbamate

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

STATEMENT OF PRACTICAL TREATMENT

148-1276

If in eyes: Flush eyes for at least 15 minutes with water. Call a physician immediately.

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION: MAY BE IRRITANT IF SWALLOWED.** Do not take internally. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is anticholinergic.

**ENVIRONMENTAL HAZARD**

This product is toxic to fish, birds and other wildlife. Do not apply directly to lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Shrimp and crab may be killed at application rates recommended on this label. Do not apply this product where these are considered important resources.

**Bee Caution:** May kill bees in substantial numbers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Entomology Service.

Do not use when value of bees as pollinators is more important than insect control. Before applying warn beekeepers to locate hives beyond bee flight range until one week after application to take other equally effective precautions.

Avoid contamination of Food, Feed, Fertilizers, and Water by leachates.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

**STORAGE AND DISPOSAL**

Do not store in a way that will contaminate the storage container or its contents.

**STORAGE:** Store in original container. May be stored in airtight plastic bags that are placed in a clean, dry, well-ventilated space. For the most accurate information, refer to the label provided at a time. Wear protective clothing and use proper disposal methods. Do not store in a way that will contaminate the storage container or its contents. Do not store in a way that will contaminate the storage container or its contents.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product should be disposed of in accordance with applicable Federal, State and local laws.

**CONTAINER DISPOSAL:** Do not reuse this container for any other purpose. If the container is to be recycled, it must be cleaned, dried, and then recycled in accordance with applicable Federal, State and local laws.

EPA Reg. No. 148-1276

EPA Est. No. 98-9111

**NET WEIGHT 50 POUNDS (22.68 kilograms)**



**THOMPSON CHEMICAL COMPANY  
CHEMICAL COMPANY**

P.O. BOX 2383, KANSAS CITY, KANSAS 66110

**BEST DOCUMENT AVAILABLE**

