

170 09
71 15 17/12
193

Anderson's®
FARM TOP
FLY SPRAY

FOR BULLY WITH BULL CALVES,
Horses, Hogs

Anderson's®
Great Fogging Spray

FARM TOP
FLY SPRAY

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response. Page 2.12, Active to Inert Change in Status.

A COPY
READY
TO USE
PRODUCT
IN
FOGGERS
AND
SPRAYERS
BY
ANDERSON

E.P.A. ESTABLISHMENT
No. 150-MN-1

E.P.A. REGISTRATION
No. 150-09



KILLS FLIES
AND MOSQUITOES

Pyrethrins .02000% - Technical Piperonyl Butoxide* .03984% -
N-octyl Bicycloheptene Dicarboximide .06667% - 2,2-Dichlorovinyl
dimethyl phosphate .4092% - related compounds .0308% - INERT
INGREDIENTS: Refined Mineral Oil 5.00000% - Petroleum Hydro
carbons 84.43368%
Equivalent to .03187% (butylcarbityl) (6-propylpiperonyl) ether and
.00797% related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Back Panel For Other Cautions




Anderson's® FARM TOP FLY SPRAY

ANOTHER PRODUCT IN THE FINE LINE OF

Anderson's® DAIRY INSECTICIDES

A READY TO USE PRODUCT



DIRECTIONS

IN BARN FOR CONTROL OF HORN FLIES, STABLE FLIES, MOSQUITOES AND GNATS: Apply at rate of 1 pint per 8,000 cubic feet. Close all doors and windows prior to spraying. Suspend spray mist in the air so it settles gradually to the floor. Keep closed for 10 to 20 minutes. Sweep up and destroy fallen insects. Repeat as necessary. Do not fog where animals have received direct applications of DDVP or Vapona within 8 hours.

FOR TREATMENT OF DAIRY AND BEEF CATTLE FOR HORN FLIES, STABLE FLIES, MOSQUITOES AND GNATS: Use a good machine sprayer or fogger and apply as a mist. We recommend 1 to 2 ounces of FARM TOP to each animal. Spray before milking or pasturing for protection from horn flies, house flies, stable flies, mosquitoes and gnats. Particular attention should be taken to cover the animal's entire body especially the legs, shoulders and belly. Avoid wetting the hides of the animals and around the eyes, nose and mouth. Do not apply within one week of slaughter. Do not exceed 2 ounces daily per animal. Care should be taken that the spray does not come in direct contact with the cow's teats unless they are washed and dried before milking. Such application may seriously degrade the quality of the milk.

FOR TREATMENT OF HORSES AND PIGS FOR STABLE FLIES, MOSQUITOES AND GNATS: Spray FARM TOP on pigs to protect them from stable flies, mosquitoes and gnats. Spray daily once or twice as needed. Spray adult hogs lightly so as to moisten the hair tips. Do not exceed 1 ounce daily per adult hog or .5 ounce for younger animals. Use mist spray only on horses and do not exceed 2 fluid oz. per animal at each application.

FOR KENNELS AND HORSE BARN TO CONTROL HORN FLIES, STABLE FLIES, MOSQUITOES AND GNATS: Apply FARM TOP at rate of 1 quart per 1,000 square feet of surface area to outside dog runways, window sills and ledges of dog kennels. Apply at rate of 1 1/2 pints per 1,000 square feet of surface area in horse barns for control of flies. Spray floors, walls and cracks where infestations may occur.

SANITATION: Breeding and travel areas like manure piles and garbage should be removed and destroyed.

CAUTION: Harmful or Fatal if swallowed, inhaled or absorbed through the skin. Do not breathe spray mist. Do not get in eyes or on clothing. In case of contact, wash immediately with soap and water. Wash all contaminated clothing with soap and hot water before re-use. Wear protective clothing. If swallowed, call a physician immediately. Wash hands, arms and face with soap and water after handling and before eating or smoking. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they show hypersensitivity to organic phosphates. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

EMPTY CONTAINER: Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

This product is toxic to fish, wildlife and birds. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risks to persons or property arising out of use or handling and accepts the product on these conditions.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. Store in original container in a secure area away from animal feed and fodder. Do not use, pour, spill or store near heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food or feed by storage or disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL: This product is toxic to fish, wildlife and birds. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Attached to label.