




*Anderson's*  
**New DDVP DRY BAIT**  
**FLY KILLER**  
 AND  
**INSECTICIDE**

*with stabilized DDUP*

"A READY-TO-USE ANDERSON INSECTICIDE  
 IN THIS HANDY SHAKER" • **KILLS RESISTANT FLIES**

ACTIVE INGREDIENTS:  
 2, 2-dichlorovinyl  
 dimethyl phosphate ----- 0.5%  
 INERT INGREDIENTS\* ----- 99.5%  
 \*Includes Bait, Sweetners

ACCEPTED



U. S. D. A.  
 REGISTRATION  
 No. 150-24

**DIRECTIONS**

**DAIRY BARNs, MILK SHEDS, STABLES AND LOAFING AREAS, LIVESTOCK BARNs, PIG PENS, AND OUTSIDE AREAS.** Scatter dry bait at rate of 1/4 lb. per 1,000 sq. ft. on floor or ground areas where flies gather. Ideal baiting sites are around doorways, feed storage rooms, alleyways, and window sills. On freshly limed floors, place bait on paper or on burlap sacks. Repeat as necessary for practical fly kill.

**CAGED POULTRY AND TURKEYS:** Spread dry bait at the rate of 1/4 lb. per 100 sq. ft. on droppings beneath cages. Repeat weekly.

**POULTRY HOUSES:** Scatter dry bait at the rate of 1/4 lb. per 100 sq. ft. where flies gather but keep out of reach of birds. Apply to manure, on window sills, outside penned enclosures and on floor of feed storage room. Repeat as necessary.

**KENNELs:** Treat walkways and other areas surrounding animal runs at the rate of 1/4 lb. per 1000 square feet. For caged units, apply to dropping area.

**GARBAGE CANS:** Sprinkle around and inside garbage cans. Repeat as needed.

Around Restaurants, Drive-Ins, Ice Cream Stands, Picnic Grounds, Backyard areas, Patios, Outdoor Latrines, sprinkle fly bait at the rate of 1/2 lb. per 1000 square feet of area on floors, walks, alleyways, and other horizontal surfaces where flies frequent.


**CAUTION:**  
 HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep out of reach of children, poultry, and domestic animals. If swallowed, induce vomiting immediately and get medical attention at once. Atropine is antidotal.

**NOTICE:**  
 Because we cannot control the application of this product by the user, the seller can assume no responsibility other than the value of this can and its contents.



**CAUTION: KEEP OUT OF THE REACH OF CHILDREN**  
 See Back Panel For Other Cautions  
 ANDERSON CHEMICAL COMPANY





*Anderson's*  
**New DDVP DRY BAIT**  
**FLY KILLER**  
 AND  
**INSECTICIDE**


*with stabilized DDVP*

A READY-TO-USE ANDERSON INSECTICIDE  
 IN THIS HANDY SHAKER • **KILLS RESISTANT FLIES**

ACTIVE INGREDIENTS  
 2, 2-dichlorovinyl  
 dimethyl phosphate 0.5%  
 INERT INGREDIENTS\* 99.5%  
 \*Includes Bait Sweeteners



**CAUTION: KEEP OUT OF THE REACH OF CHILDREN**  
 See Back Panel For Other Cautions  
**ANDERSON CHEMICAL COMPANY**



ACCEPTED

U. S. D. A.  
 REGISTRATION  
 No. 150-24



**DAIRY BARNs, MILK SHEDS, STABLES AND LOAFING AREAS, LIVESTOCK BARNs, PIG PENS, AND OUTSIDE AREAS.** Scatter dry bait at rate of 1/4 lb. per 1,000 sq. ft. on floor or ground areas where flies gather. Ideal baiting sites are around doorways, feed storage rooms, alleyways, and window sills. On freshly limed floors, place bait on paper or on burlap sacks. Repeat as necessary for practical fly kill.

**CAGED POULTRY AND TURKEYS:** Spread dry bait at the rate of 1/4 lb. per 100 sq. ft. on droppings beneath cages. Repeat weekly.

**POULTRY HOUSES:** Scatter dry bait at the rate of 1/4 lb. per 100 sq. ft. where flies gather but keep out of reach of birds. Apply to manure, on window sills, outside penned enclosures and on floor of feed storage room. Repeat as necessary.

**KENNELs:** Treat walkways and other areas surrounding animal runs at the rate of 1/4 lb. per 1000 square feet. For caged units, apply to dropping area.

**GARBAGE CANS:** Sprinkle around and inside garbage cans. Repeat as needed.

Around Restaurants, Drive-Ins, Ice Cream Stands, Picnic Grounds, Backyard areas, Patios, Outdoor Latrines, sprinkle fly bait at the rate of 1/2 lb. per 1000 square feet of area on floors, walks, alleyways, and other horizontal surfaces where flies frequent.

**CAUTION:**  
 HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep out of reach of children, poultry, and domestic animals. If swallowed, induce vomiting immediately and get medical attention at once. Atropine is antidotal.

**NOTICE:**  
 Because we cannot control the application of this product by the user, the seller assumes no responsibility other than the value of this can and its contents.



NET CONTENTS 1 LB.