



CIRECTIONS

IN BARN: Close all doors and windows prior to mitking. Suspend spray mist in the air so it settles gradually to the floor. Keep closed for 10 to 20 minutes. Sweep up and destroy fallen insects. Repeat as necessary for control of flies, mosquitos and gnats. Apply at rate of 1 pint per 8,000 cubic feet.

HORN FLIES, STABLE FLIES, MOSQUITOS AND GNATS: For dairy and beef cattle, horses and hogs, use a good hand machine sprayer or fogger and apply as a mist. We recommend 1 or 2 ounces of FLYGA KVIST to each animal. Spray before milking or pasturing for protection from horn flies, house files, atable files, mosquitos and gnats. Particular attention should be taken to cover the animal's entire body, especially the legs, shoulders and belly. Avoid wetting the hides of the animals and around the eyes, nose and mouth. Do not apply within one week of slaughter. Do no exceed 2 ounces daily per animal. Care should be taken that the spray does not come in direct contact with the lactating dairy cow's tests unless they are washed and dried before milking. Such application may seriously degrade the quality of the milk.

IN BARN FOGGER: To kill horn files, stable files, barn files, mosquitos and gnats, apply at the rate of 1 pint per 8,000 cubic feet. Before spraying reduce air movement by closing doors, windows and other openings. Do not fog where animals have received direct application of DDVP within 8 hours.

SANITATION: FLYGA KVIST will be more effective when breeding and travel areas like manure piles, garbage disposals, etc. are removed and destroyed.

WARNING: Harmful or Fatal If swallowed, Inhaled or absorbed through the skin. Do not breathe spray mist. Do not get in eyes or on clothing. In case of contact, wash immediately with soap and water. Wash all contaminated clothing with soap and hot water before reuse. Wear protective clothing. If swallowed, induce vomiting immediately and get medical attention. Do not contaminate water, feed, milk or milking equipment. Do not store in temperatures over 130 degrees F. Keep well corked when not in use. Do not apply regularly to calves under 6 months of age. Braham cattle should not be treated as they may show hypersensitivity to organic phosphates. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. Store in original container in a secure area away from animal feed and fodder. Do not use, pour, spill or store near heat or open flame.

PESTICIDE DISCOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not conteminate water, food or feed by storage or disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL: This product is toxic to fish, wildlife and birds. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NOTICE

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risks to persons or property arising out of use or handling and accepts the product on these conditions.

MANUFACTURED BY ANDERSON CHEMICAL COMPANY LITCHFIELD, MINNESOTA 55355