

## DIRECTIONS FOR USE

**INDOOR APPLICATION:** The use of this product in food processing establishments should be restricted to those periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended. Vacate treated area and ventilate before reoccupying. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

**SURFACE SPRAYING:** Treat all accessible exposed stages of roaches, ants, silverfish, spider mites, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-tooth grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites, and cadelles. Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at the rate of 1 gallon to 750 square feet of surface. To control carpet beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs, spray mattresses lightly, particularly around tufts and seams. Take bed to attic and spray in all joints. Treat baseboards, mouldings, and floor. Repeat treatment as needed.

To control lice and tracheomonas, thoroughly spray infested animal quarters, resting quarters, nearby cracks and crevices, lining and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths, remove any infested articles from storage, wash thoroughly and hang for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

# PIONEER AQUEOUS SUPER-CIDAL INSECTICID

## A GENERAL PURPOSE AQUEOUS INSECTICIDE

A Clear, Stable, Micro Emulsion - Leaves No Objectional Residue.

Effective fast acting insecticide for use in homes, institutions, restaurants, food processing plants, dairies, outdoors for Mosquito control. Apply with conventional mechanical or compressed air equipment.

### ACTIVE INGREDIENTS:

Pyrethrins	0.1
*Piperonyl Butoxide, technical	1.0
Petroleum distillate	0.9

### INERT INGREDIENTS:

18 equivalent to 0.8% (butylarbityl) for total inert ingredients in formulated compounds 98.0

EPA Reg. No. 151-16-AA

MANUFACTURED AND DISTRIBUTED BY



"PIONEERS IN SANITATION"

2051 VIOLET STREET, LOS ANGELES, CALIF. 90024

PHONE 624-2869

**CAUTION:** KEEP OUT OF REACH OF CHILDREN  
RIGHT PANEL FOR ADDITIONAL CAUTION

NET CONTENTS 1 GALLON