

ACCEPTED

AUG 3 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REG. STER-
ED UNDER NO. 168-59 SUBJECT
TO ATTACHED COMMENTS.

**WASATCH CHEMICAL COMPANY
SALT LAKE CITY, UTAH**

CENTER PANEL

PARATHION 25%

EMULSIBLE

Contains 2 Pounds of Parathion Per Gallon

**DANGER:
KEEP OUT OF REACH OF CHILDREN
READ ALL DIRECTIONS CAREFULLY
BEFORE USING!**

ACTIVE INGREDIENTS:

Parathion (O,O-Diethyl O-p-nitrophenyl thiophosphate).... 25.0%
Aromatic petroleum derivative solvent..... 70.0%
INERT INGREDIENTS:..... 5.0%

TOTAL.....100.0%

 **P O I S O N**

U.S.D.A. Reg. No. 168-59

CONTENTS: 5 GALLONS

Manufactured by:
WASATCH CHEMICAL COMPANY
SALT LAKE CITY, UTAH

RIGHT PANEL -

DIRECTIONS FOR USE

Carefully read all directions before using this material.

VEGETABLE PESTS: Apply 1 to 2 pints per acre in enough water to give a thorough and complete coverage. Make initial application when pests first appear in numbers, and repeat as necessary. Consult State Agricultural Extension Service for proper timing of application and exact dosage.

For control of aphids, army worms, cabbage worms, celery worms, Colorado potato beetle, cucumber beetles, flea beetles, grasshoppers, leafhoppers, leafminers, melon worms, pea weevil, pickle worms, stink bugs, thrips, vine borer, webworms, white fly on beans, beets, broccoli, cabbage, carrots, cauliflower, celery, corn, cucumbers, lettuce, melons, onions, peas, potatoes, pumpkins, spinach, squash, tomatoes. Do not use later than 5 days before harvest on potatoes; 7 days on broccoli, cauliflower; 10 days on cabbage, peas, pumpkins, tomatoes; 12 days on corn; 14 days on spinach; 15 days on beans, beets, carrots, cucumbers, melons, onions, sweet potatoes, squash; 21 days on beet tops for celery, greens and lettuce. Do not apply Parathion 25% Emulsible to cucumbers before vining.

CROP PESTS: Apply 1 to 2 pints per acre in enough water to give thorough and complete coverage. Make initial application when pests first appear in numbers, and repeat as necessary.

For control of alfalfa caterpillar, aphids, army worms, grasshoppers, lygids, bugs and mites on alfalfa, clover, and small grains (wheat, oat, barley). Do not apply later than 15 days before harvest on crops listed above.

FRUIT PESTS: Mix 1 to 2 pints per 100 gallons of water and apply when pests first appear in numbers and repeat as necessary. Do not apply more than 2 1/2 gallons of Parathion 25% Emulsible per acre to peaches. Use on peach areas other than California. On apples and pears for control of aphids, codling moth, mites, leafhoppers, slugs, San Jose scale and crawlers, grasshoppers, shot-hole borers, woolly aphids; on cherries for aphids, mites, pear slug, San Jose scale crawlers; on peaches for lygus bugs, shot-hole borer, red hant leafroller; on prums-

LEFT PANEL -

and prunes for aphids, mites, leafhoppers; on raspberries for mites; on strawberries for weeviler and mites.

Do not use more than 500 gallons of dilute spray per acre on peaches, pears and prunes, 700 on pears or 400 on cherries.

Do not apply later than 14 days before harvest on apples, pears, cherries, peaches, plums, prunes, strawberries. On raspberries apply only before bloom or after harvest.

Parathion may cause injury to fruit and foliage on McIntosh and related varieties of apples.

ORNAMENTAL PESTS: For commercial plantings only, not around dwellings.

For control of aphids, white flies, mites, leafhoppers, thrips, lace bugs, bag worms, cottony cushion scale, lecanium scale, mealy bugs, and leaf-tiers.

Mix 1 to 2 pints per 100 gallons of water and apply when pests first appear in numbers and repeat as necessary.

DANGER: Poisonous if swallowed, inhaled, or absorbed through skin; rapidly absorbed through skin! Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection.

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate food and foodstuffs. Wash hands, arms, face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds.

Apply this product only as specified on this label.

Do not reuse empty container. Destroy by crushing or perforating and ^{with care} ~~disposing~~.

ANTIDOTE AND FIRST AID TREATMENT

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING.

SWALLOWED, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet.

IF ON SKIN: In case of skin contact, remove contaminated clothing and immediately wash skin with soap and water.

WARNING: Be sure to read and understand the precautionary statements before using. Do not prune trees, thin fruit, or work in dense orchards for two weeks after spraying to allow vapors to dissipate. Keep containers closed when not in use, and do not re-use for other purposes. Do not store near food or feed materials. Destroy container when empty. Combustible. Do not store or use near heat or open flame.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label.