

CHLORDANE DUST

Ready to use, apply as directed.

Controls: Ants - Box Elder bugs - Cockroaches - Cutworms - Earwigs - Grasshoppers - Grubs in Lawns - Maggots - Silverfish - Soldiers - Water bugs, and others listed below.

ACTIVE INGREDIENTS:

Technical Chlordane ..... %  
Equivalent to 3% octachlorocyclohexane, 1-methoxy-1-trifluoromethyl-2,2,4,4-tetrahydro-1H-benzopyrene and 2% of related compounds.

INERT INGREDIENTS:

..... 95%  
TOTAL 100%

5-1-67

168-98  
comments

CAUTION: KEEP OUT OF REACH OF CHILDREN.

U.S.D.A. Reg. No. 156-96

DIRECTIONS

Chlordane kills insects by contact. Many hours are often required before insect dies. Apply dust evenly over the area to be treated. One ounce of dust covers 75-100 square feet of surface. Repeat treatment as necessary. Residues kill for several days.

ANTS: Dust over hills and along runways inhabited by ants and drive the dust into the openings of the hills if these can be located. Dust around door sills, window frames, water pipes, baseboards, etc. where ants may enter buildings, and other points where they may enter buildings. Dust directly on ants whenever possible.

BOX ELDER BUGS: Apply insecticide over soil, debris, tree trunks, foundations, etc., when and where bugs congregate. Cover door sills, window frames.

COCKROACHES, CRICKETS, SILVERFISH, WATER BUGS: Drive dust into cracks and crevices around baseboards, water pipes, window and door sills, base and under appliances, and other dark places where insects hide.

CUTWORMS: Cover entire soil area around young transplants such as annual flowers, tomatoes, cabbage and cauliflower with dust as early as possible when cutworms cause damage. Avoid contamination of edible portions or remove residues at harvest.

EARWIGS AND MILLETS: Water areas to be treated to roughly the level before treatment is to be applied. Apply dust under shrubs, flowers, etc., and over mulch, leaves and debris where earwigs hide. Avoid disturbing the treated area by watering or cultivation for a few days after treatment.

GRASSHOPPERS: Treat green plants such as corn, sorghum, and weeds in waste areas, preferably when grasshoppers are young and before they migrate to cultivated plants, or later on green plants for larger grasshoppers. Do not allow livestock to graze on treated areas.

GRUBS, EARTHWORMS, ANNECED BEETLES, AND MILLIPEDS IN LAWN: Apply dust at rate of 100 lbs. per 100 square feet of lawn (10 lbs. per 10 ft. by 10 ft. in single garden or later when damage begins to appear. Wash dust off soil with water after treatment. Avoid contamination of edible portions.

CRICKETS, CHINCHIPES, AND MILLIPEDS: Apply dust at rate of 100 lbs. per 100 sq. ft. of lawn. In single garden or later when damage begins to appear. Wash dust off soil with water after treatment. Avoid contamination of edible portions.

STRAWBERRY: Apply dust at rate of 100 lbs. per 100 sq. ft. of lawn. In single garden or later when damage begins to appear. Wash dust off soil with water after treatment. Avoid contamination of edible portions.

ONION, RADISH AND BEAN SEEDLINGS: Apply dust at rate of 100 lbs. per 100 sq. ft. of lawn. In single garden or later when damage begins to appear. Wash dust off soil with water after treatment. Avoid contamination of edible portions.

SPIDERS: Apply Chlordane dust to areas where spiders are found. In corners or wherever spiders may be found. Repeat as necessary.

CAUTION: Avoid inhalation of dust. Avoid contact with eyes.

This product is toxic to fish and aquatic life. Do not apply to water. Apply this product as directed.

NOTICE: Manufacturers of pesticides should not use this product in their formulations. Do not use this product in formulations.

MADE IN U.S.A.

REGISTERED TRADEMARK