

WASATCH CHEMICAL COMPANY
SALT LAKE CITY, UTAH

ACCEPTED
FEB 14 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 168-171 SUBJECT
TO ATTACHED COMMENTS.

PREPARATION

ACTIVE INGREDIENTS:
100% DUST
100-b ethyl (2-pyridyl) phosphorothioate..... 98%
INERT INGREDIENTS..... 2%
Total..... 100%

DANGER: KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANELS FOR OTHER CAUTIONS.

O P O I S O N O
X X X X X X

U. S. D. A. REG. NO. 168-171

DIRECTIONS

VEGETABLE CROPS:

For control of aphids and mites on beans, beets, carrots, cabbage, celery, pepper, eggplant, cucumber, squash, melon, lettuce, peas, potatoes, tomatoes and spinach, apply 10 to 20 pounds of dust per acre when pest appears in numbers and later as necessary. Direct dust to underside of leaves for mite control. The higher rate is for heavier foliage. Do not treat beets within 21 days of harvest if tops are to be used for food or feed.

On beans for control of Mexican bean beetle, leafhoppers, aphids, red spiders, leaf rollers, armyworms, and leaf miners, apply 15 to 20 pounds of dust per acre and repeat in 7 to 10 days if necessary.

For control of leafhoppers on lettuce; Colorado potato beetle, flea beetle, leafhopper, sculptured leaf miner, grasshoppers, aphids, and white fly on potatoes, red tomatoes, midwest thrips on onions; grasshoppers, armyworms, and European corn borer on corn, apply 15 to 20 pounds of dust per acre and repeat as necessary.

Do not apply within 14 days prior to harvest to lima beans, dry beans, beets, carrots, eggplant, melon, squash, peppers and peas and within 7 days of harvest to snap beans, cucumbers, broccoli, broccoli sprouts, celery, celery, cucumbers, onions, potatoes, tomatoes, tomato, tomato and tomato within 11 days of picking harvest as a crop.

CRUITS:

On alfalfa and flower for control of aphids, armyworms, blister beetles and grasshoppers and on small grains for control of aphids and webworms; apply 15 to 20 pounds of dust per acre when pest first appears in numbers and later as necessary. Make no applications within 7 days of cutting for hay or feed.

ESCAL CROPS:

On alfalfa and flower for control of aphids, armyworms, blister beetles and grasshoppers and on small grains for control of aphids and webworms; apply 15 to 20 pounds of dust per acre when pest first appears in numbers and later as necessary. Make no applications within 7 days of cutting for hay or feed.

PARATHION (85%)

WARNING:

May be fatal if swallowed, inhaled, or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes or on skin or clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unpracticed persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

ANTIDOTE:

INTERNAL: Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep warm. Call a physician.

EXTERNAL: If skin is accidentally contaminated, wash thoroughly with soap and warm water.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Do not apply when weather conditions favor drift of dust from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (Keep out of tanks).

NON-WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product. Buyer assumes all responsibility for handling whether in a lot or as individual containers.

PARATHION (85%)

BERNARD CHEMICAL COMPANY
NEW YORK, N. Y.