

ACCEPTED

10-1-1916

U.S. DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY
WASHINGTON, D. C.

APPLICANT'S NAME:
ADDRESS:
CITY:
STATE:
COUNTRY:
DATE OF RECEIPT:
AMOUNT PAID:
TOTAL: 175.00

COMMENTS:
Aphids, Blister Mite, Leafhoppers, Red Spider Mite, Rose Slug, Scale, Crawlers, Black Leaf Curl, Powdery Mildew, Shot Hole Fungus, Rust.
168-117

For control of peach leaf curl, shot hole fungus, powdery mildew, and aphids, blister mite and scale crawlers on apples, pears and stone fruits, mix 1 teaspoon per gallon of water and apply it when buds begin to swell in spring; again at pink bud stage and later at petal fall stage.

To control brown rot and blight on citrus, spray during winter months before picking. For control of ash blight, spider, exposed thrips and scale crawlers, spray when pest appears in number 10 to 15 days later if necessary for control.

For control of aphids, red spider, rose slug, leafhoppers, dipterous beetles, and thrips, plus powdery mildew, leaf spot, rust, black spot on such plants as roses, carnations, dahlias, gladioli, sweet peas, use 4 teaspoons per gallon, and apply it when insects or diseases first appear, and later at weekly intervals if necessary for control.

For control of aphids, red spider mite, leaf roller, and leaf spot on strawberries and raspberries, mix 1 teaspoon per gallon, and apply when pest first appears, and 10 to 15 days later if needed.

Harmful if swallowed. Avoid contact with solutions and spray plants. Avoid contamination of food and furniture. Keep out of reach of children. Wash out sprayer after use. Do not apply to fruits less than 1 day before harvest.

U.S. DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY
WASHINGTON, D. C.
1916