

DIAZINON 2% DUST

USE
FOR L
ED UNDER S. 166-254
TO ATTACHED COM. 112

ACTIVE INGREDIENTS:	O,O-Diethyl O-(2-Isopropyl-4-Methyl-6-Pyrimidinyl)	
	Phosphorothiate.....	2%
INERT INGREDIENTS:	98%
		TOTAL 100%

DIRECTIONS

For control of aphids, leaf miners, thrips, leafhoppers, on table beets, broccoli, cabbage, cauliflower, canteloupes, cucumbers, squash, watermelons, peppers, potatoes, beans, tomatoes, turnips, and strawberries. Apply 20 to 35 lbs. per acre when pest appears in number, and repeat as needed. Do not apply to tomatoes within three days of harvest; peppers and strawberries 5 days; beans, broccoli, cabbage, cauliflower, canteloupes, cucumbers, squash and watermelons 7 days; turnips 10 days; beets and potatoes 14 days before harvest. Do not exceed 0.5 pound of actual Diazinon per acre on beets.

COCKROACHES: For control of cockroaches inside and under dwellings, dust thoroughly the areas around water pipes, cracks, and crevices; behind and beneath cabinets, refrigerators, sinks and other places where cockroaches crawl or hide. Retreat as necessary. Diazinon dust in dwellings is limited to pest control operators only.

CAUTION: Harmful if swallowed.
Avoid prolonged breathing of dust.
May be absorbed through the skin.
Avoid contact with skin and eyes.
Avoid contamination of feed and foodstuffs.
Wash thoroughly after handling.
Keep out of reach of children.

NON-WARRANTY: Seller makes no warranty of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risks of use or handling, whether in accordance with directions or not.

NET CONTENTS:

MANUFACTURED BY: MASATCH CHEMICAL COMPANY
SALT LAKE CITY, UTAH

May 18, 1962

S. J. ...