

Contains 0.5% ^RCORAL

For Control of Certain Ectoparasites of Animals

ACTIVE INGREDIENTS:

<u>O,O</u> -Diethyl <u>O</u> -3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl phosphorothioate	- 0.5%
INERT INGREDIENTS.....	- 99.5%
Total.....	100.0%

U. S. Patent No. 2,748,146
USDA Reg. No. 168-299 Rev.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE REAR PANEL FOR OTHER CAUTION STATEMENTS.

STOP: READ LABEL BEFORE USE

Directions for Use

Parasite Dust is suitable for direct application by any conventional dusting equipment such as rotary or hand dusters, puff dusters, or by sprinkling from a can.

Parasite Dust is an effective insecticide for the control of certain wool, hair, and flesh parasites. When these parasites are abundant, the animals are unthrifty and unprofitable.

Recommended Applications

<u>Animal</u>	<u>Parasite</u>	<u>Remarks</u>
<u>Poultry</u>	Northern Fowl Mite	Direct Application: Dust birds thoroughly, especially under wings and around vent. Supplemental control can be obtained by also treating litter and housing. Repeat as necessary but not more often than weekly.
	Northern Fowl Mite Poultry Red Mite Lice Flies	Litter & Housing Treatment: Apply 1.4 lbs. per 100 sq. ft. of surface area to walls, ceiling, floor, <u>roosts, nests and adjacent areas</u> , and 1 lb. per 20 sq. ft. of litter area. Force dust into all cracks and crevices. Repeat as necessary.
<u>Sheep</u>	Sheep Ked (ticks) Fleece Worms Wool maggots	Hand Duster: Apply 1 to 2 ozs. of dust per head. Dust on and rub into wool over entire body. Power Duster: Machine setting 1½ to 4 lbs. of dust per minute. Treatment Rate: 50 to 100 sheep per minute. Repeat as necessary but not within 15 days of slaughter.

11/3/66

ACCEPTED

FEB 21 1967

UNDER THE FEDERAL INSECT
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 168-299 SUBJECT
TO ATTACHED COMMENTS

Animal	Parasite	XXXXXXXXXXXXXXXXXXXX	Remarks
<u>Dogs</u>	Fleas Lice Ticks	XXXXXXXXXXXX	Dust dog thoroughly under the hair, over entire body. Repeat as necessary, but not more than weekly. Dog's bedding should also be dusted to help prevent reinfestation.

Caution

For external insecticidal use on animals.
 Keep out of the reach of children.
 Avoid prolonged inhalation of dust.
 Wash after handling.
 Do not contaminate feed or food.
 If misuse results in accidental poisoning, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. Atropine sulfate is antidotal.

Restrictions

Do not use on cats or nursing puppies. Do not treat poultry within 10 days of vaccination or other stress influence. Avoid contamination of feed, water, and water utensils. Provide thorough ventilation while dusting.

NON-WARRANTY: Wasatch Chemical Company makes no warranty of any kind, expressed or implied, concerning the use of this material, except that it conforms to the chemical description on the label. Neither Wasatch Chemical Company or the Seller shall be held responsible in any manner for any personal injury or property damage, or other type of loss resulting from the handling, storage, or use of this material. The Buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

NET CONTENTS: 50 lbs.

^RCORAL is Registered Trade Mark of Farbentabriken Bayer, A. G., Chemagro Corp., licensee.

3-30-66