

\$2.95




ROSE FOOD

PLUS SYSTEMIC INSECTICIDE

16-16-8

CONTAINS 1% DI-SYSTON®*

Protection for
ROSES • BULBS • SHRUBS • FLOWERS



CONTROLS Aphids, Birch Leaf Miner, Lace Bug, Leafhopper, Mites, Thrips, White Flies, Mimosa Webworm and Pine Tip Moth.

ROSE FOOD PLUS SYSTEMIC INSECTICIDE

16-16-8

GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N)	16.00%
10.7% Ammoniacal Nitrogen from Ammonium Nitrate Sulfate, Monoammonium Phosphate and Ammonium Chloride	
5.3% Nitric Nitrogen from Ammonium Nitrate Sulfate and Potassium Nitrate	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	16.00%
Derived from Monoammonium Phosphate, Fluor-apatite and Hydrated Phosphate of Iron and Aluminum	
SOLUBLE POTASH (K ₂ O)	8.00%
Derived from Potassium Chloride and Potassium Nitrate	

ROSE FOOD PLUS SYSTEMIC INSECTICIDE enters the plant by absorption through the roots. The insecticide placed near the roots is translocated to all parts of the plant, killing sucking insects feeding on the plant.

FOR BEST RESULTS — USE THE MEASURING CUP
Level-full equals 2 oz., enough for one rose bush.

DIRECTIONS FOR USE

- ROSES:** Use 1 measure per hybrid tea bush 18 to 24 inches in height around drip area of plant. Water normally after.
- CLIMBERS:** Apply 1 measure per foot of cane in early spring. Repeat before July 15th if pests reappear or if feeding is needed.
- NEW PLANTINGS:** Mix 1 measure in bottom of hole before planting. Water normally after planting.

SHRUBS & EVERGREENS

Amount to Use	Height of Plant
2 oz. per ft. of height	1-3 feet: 1 measure full
4 oz. per ft. of height	3-6 feet: 2 measures full
6 oz. per ft. of height	6-12 feet: 3 measures full

Use a shovel or bar and place insecticide-fertilizer 4 to 6 inches underground within drip area of plant. Water normally after application. Apply at two or more places per plant.

FLOWERS AND BULBS: (Dahlias, Iris Corms)


Broadcast 5 lbs. per 300 sq. ft. or 1 lb. per 60 sq. ft. of area. Work into soil before planting. Water normally after planting. Repeat before July 15th if feeding is needed.

5 LBS. TREATS 40 Rose Bushes or Shrubs 18 to 24 inches in height, or 300 sq. ft. of flower area.

Dec. 1, 1974

168-334

ROSES • BULBS • SHRUBS • FLOWERS



CONTROLS Aphids, Birch Leaf Miner, Lace Bug, Leafhopper, Mites, Thrips, White Flies, Mimosa Webworm and Pine Tip Moth.

THE ACTIVE INGREDIENT IS AN ORGANIC PHOSPHATE INSECTICIDE AND WILL INHIBIT CHOLINESTERASE.

ACTIVE INGREDIENT:

*0,0-diethyl S-[2-(ethylthio)ethyl] phosphorodithioate 1.0%

INERT INGREDIENTS: 99.0%

(Relative to insecticide content) **TOTAL** 100.0%

*DI-SYSTON — Registered trademark of Farbenfabriken-Bayer, Chemagro Corporation License. U.S. Patent No. 2,759,010.

WARNING:

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR OTHER WARNINGS

EPA Reg. No. 169-334

NET CONTENTS: 5 LBS.



around drip area of plant. Water normally after.
CLIMBERS: Apply 1 measure per foot of cane in early spring. Repeat before July 15th if pests reappear, or if feeding is needed.
NEW PLANTINGS: Mix 1 measure in bottom of hole before planting. Water normally after planting.

SHRUBS & EVERGREENS

Amount to Use	Height of Plant
2 oz. per ft. of height	1-3 feet 1 measure full
4 oz. per ft. of height	3-6 feet 2 measures full
6 oz. per ft. of height	6-12 feet 3 measures full

Use a shovel or bar and place insecticide-fertilizer 4 to 6 inches underground within drip area of plant. Water normally after application. Apply at two or more places per plant.

FLOWERS AND BULBS: (Dahlias, Iris Corms)

Broadcast 5 lbs. per 300 sq. ft. or 1 lb. per 60 sq. ft. of area. Work into soil before planting. Water normally after planting. Repeat before July 15th if feeding is needed.

5 LBS. TREATS 40 Rose Bushes or Shrubs 18 to 24 inches in height, or 300 sq. ft. of flower area.

WARNINGS

Poisoning may occur as a result of exposure. **MAY BE FATAL IF SWALLOWED.** If swallowed, induce vomiting and call physician. May be absorbed through skin. Do not get on skin, in eyes or on clothing. Wear rubber gloves. Wash hands thoroughly after handling. **KEEP OUT OF REACH OF CHILDREN.** Keep children and pets off treated area until insecticide has been washed into soil and treated areas have dried completely.


DO NOT APPLY TO PLANTS TO BE USED FOR ANIMAL OR HUMAN CONSUMPTION. Do not apply broadcast to lawn or turf areas. This product is **TOXIC** to fish, wildlife and birds. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. **DO NOT REUSE CONTAINER.** Crush and dispose of when empty. Apply this product only as specified on label.

TO PHYSICIAN — ANTIDOTE

Antidote is atropine sulphate. 2-PAM is also antidotal and may be administered in conjunction with atropine.

NON-WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purpose indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.

Protection for
ROSES • BULBS • SHRUBS • FLOWERS



CONTROLS Aphids, Birch Leaf Miner, Lace Bug,
 Leafhopper, Mites, Thrips, White Flies,
 Mimosa Webworm and Pine Tip Moth.

THE ACTIVE INGREDIENT IS AN ORGANIC PHOSPHATE INSECTI-
 CIDE AND WILL INHIBIT CHOLINESTERASE.

ACTIVE INGREDIENT:

0,0-diethyl S-[2-(ethylthio)ethyl] phosphorodithioate 1.0%

INERT INGREDIENTS:

(Relative to insecticide content) TOTAL 99.0%

DI-SYSTON -- Registered trademark of Farbenfabriken-
 Bayer, Chemagro Corporation License. U.S. Patent
 No. 2,759,010

WARNING:

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR OTHER WARNINGS

EPA Reg. No. 168-334

NET CONTENTS: 5 LBS.

ROSES

CLIMBERS

NEW PLANTINGS

FLOWERS AND SHRUBS

5 LBS. TREATS 1000 SQ. FT. OF FLOWER AREA

WARNINGS

Poisoning may occur as a result of exposure.
MAY BE FATAL IF SWALLOWED. If swallowed, induce
 vomiting and call physician.
 May be absorbed through skin. Do not get on skin,
 in eyes or on clothing. Wear rubber gloves. Wash
 hands thoroughly after handling.
KEEP OUT OF REACH OF CHILDREN. Keep children
 and pets off treated area until insecticide has
 been washed into soil and treated areas have
 dried completely.

**DO NOT APPLY TO PLANTS TO BE USED FOR ANIMAL
 OR HUMAN CONSUMPTION.** Do not apply broad-
 cast to lawn or turf areas.
 This product is **TOXIC** to fish, wildlife and birds.
 Keep out of lakes, streams or ponds. Do not
 contaminate water by cleaning of equipment or
 disposal of wastes.
DO NOT REUSE CONTAINER. Crush and dispose of
 when empty.
 Apply this product only as specified on label.

TO PHYSICIAN - ANTIDOTE

Antidote is atropine sulphate. 2-PAM is also antidotal and may be administered in con-
 junction with atropine.

EI



DISTRIBUTED BY
WASATCH CHEMICAL DIVISION

1979 SOUTH SIXTH WEST P.O. BOX 6219 SALT LAKE CITY UTAH 84106