

(FRONT PANEL)

ACCEPTED

FEB 15 1967

UNDER THE INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC FOREST REGISTERED  
UNDER NO. 14-1334 SUBJECT  
TO ATTACHED COMMENTS.

70% AVAILABLE CHLORINE

ACTIVE INGREDIENTS: CALCIUM HYPOCHLORITE..... 70%

INERT INGREDIENTS: TABLETS IN PLASTIC POUCH..... 30%

WARNING: KEEP OUT OF THE REACH OF CHILDREN.  
SEE SIDE PANELS FOR OTHER PRECAUTIONS.

X O X  
O POISON O  
X X

WASATCH CHEMICAL COMPANY  
SALT LAKE CITY, UTAH  
U.S.D.A. REG. NO. 168

(SIDE PANEL)

ANTIDOTE:

EXTERNAL: Wash with water.

INTERNAL: Drink milk, rice gruel, ouillage or raw egg white.

Follow with emetic (tablespoon of mustard in glass of water).

CALL PHYSICIAN.

CAUTION:

Close container tightly after using.

Store in cool, dry place.

Keep away from heat, acids, and organic or combustible materials.

Do not handle with wet hands.

Wash hands immediately after handling.

Avoid prolonged contact with skin. Do not get in eyes.

Do not take internally. May produce chemical burns. In case of eye contact, flush with plenty of water. If irritation persists, get medical attention.

(RIGHT PANEL)

DIRECTIONS FOR USE

Swimming Pool Sanitation: Use Wasactor Tablets by suspending polyethylene net bags or baskets containing the proper amount of tablets about two feet below the water level in the pool. There should be more tablets in the deep end. In freshly filled pools or pools that have not been previously treated, Repeat the daily dosage until a chlorine residual of 0.6 to 1.0 ppm is obtained.

Net contents:

WASATCH CHEMICAL COMPANY  
SALT LAKE CITY, UTAH