

(FRONT PANEL)

WASAFLOR GRANULAR

Granular Calcium Hypochlorite
70% Available Chlorine

ACTIVE INGREDIENTS:	Calcium Hypochlorite.....	70%
INERT INGREDIENTS:	Inert fillers, etc.....	30%
	TOTAL	100%

DANGER: KEEP OUT OF THE REACH OF CHILDREN.
SEE SIDE PANELS FOR OTHER PRECAUTIONS

X		X
O	P O I S O N	O
X		X

U. S. D. A. REG. NO. 168-337

(LEFT PANEL)

ANTIDOTE:

EXTERNAL: Wash with water.
INTERNAL: Drink milk, rice gruel, mucilage or raw egg white.
Follow with emetic (tablespoon of mustard in glass of water). Call Physician.

CAUTION:

May be fatal if swallowed.
Close container tightly after using.
Store in a cool, dry place.
Keep away from heat, acids, and organic or combustible materials.
Do not handle with wet hands.
Do not get in eyes or on skin.
Wash hands immediately after handling.
Avoid prolonged contact with skin.
Do not take internally. May produce ^{severe} chemical burns. In case of eye contact, flush with plenty of water. If irritation persists, get medical attention.

~~.....~~
~~.....~~
~~.....~~

(RIGHT PANEL)

DIRECTIONS FOR USE

Swimming Pool Sanitation: Maintain a chlorine residual level of 0.6 to 1.0 parts per million (ppm) ~~.....~~. Available chlorine is determined with a kit which can be obtained from your swimming pool supply dealer. For initial treatment, use 1 oz. of Wasaflor Granular for every 5000 gallons of water, or two quarts of stock solution per 5000 gallons of water. (See table for stock solution preparation). Wait 30 minutes and

RECEIVED
FEB 6 1967
U. S. D. A. REG. NO. 168-337

(RIGHT PANEL)

check chlorine residual with test kit. Repeat the initial treatment if necessary. Sunlight, temperature, bathing load, and air-borne contaminants will affect the chlorine residual level so the pool water must be checked several times daily and adjustments made to maintain the level of residual chlorine. To maintain the chlorine residual level, use approximately 1/2 oz. (about 1 tablespoonful) of Wasaclor Granular per 5000 gallons of pool water or 1-1/5 quarts of stock solution.

Wasaclor Granular Needed to Provide An
Equivalent of 0.6 PPM Available Chlorine

Gallons of Water	Wasaclor Granular		Stock Solution
	Ounces	Tablespoons	
500	0.06	0.1	1/4 pint
1,000	0.12	0.2	1/2 pint
5,000	0.60	1.0	1 1/4 quarts
10,000	1.2	2.0	2 1/2 quarts
15,000	1.8	3.0	3-3/4 qts.
20,000	2.4	4.0	5 qts.
25,000	3.0	5.0	6 1/4 qts.
100,000	12.0	20.0	25.0 qts.

Preparation of Stock Solution

Use a crock or other non-metallic vessel for preparing and storing stock solution.

Wasaclor Granular

1 1/4 pounds
2 1/2 pounds
5 pounds

Lukewarm Water

10 gallons
20 gallons
40 gallons

Alkalinity or pH must be kept between 7.2 and 7.4 to minimize eye and skin irritation. A daily check with a test kit should be made. Consult your swimming pool supply dealer for methods of adjusting pH.

Super chlorination and algae control: To control algae, increase the Wasaclor Granular dosage 5 to 10 times the usual amount. This should be done once each week, preferably at night, when the pool is not in use. To keep your pool clean and help prevent algae, backwash the filters frequently, use a pool vacuum cleaner and clean the surface with a skimmer. Super chlorination also precipitates metallic impurities which may be removed with the pool vacuum.

NET CONTENTS:

MANUFACTURED BY: WASATCH CHEMICAL COMPANY
SALT LAKE CITY, UTAH

November 17, 1965