


WASCO
44-0-0 (B)
Contains 1.0% DI-SYSTON

Active Ingredients:

Di-syston 1.0% []



Manufactured by...
Net Contents:
EPA Reg. No. 166-

DIRECTIONS FOR USE

DI-SYSTON is a systemic insecticide that requires use only 100 to 200 lbs.

The insecticide, when placed near the roots, is picked up and moves in the sap stream to all parts of the plant where it kills insects as they feed on the plant.

Beans: For control of aphids, leafhoppers, Mexican bean beetle, plant bugs, mites and rips. Apply 100 to 200 lbs. (1 to 2 lbs. actual DI-SYSTON) per acre. For snap or green bean beans, place WASCO 44-0-0 in a band on each side of the seed furrow at planting. For the snap and green bean beans, place WASCO 44-0-0 in a band on each side of the seed furrow at planting. For established beans, apply 100 to 200 lbs. per acre in a band on each side of the seed furrow at planting.

Water: When used in water, apply at the rate of 100 to 200 lbs. per acre under conditions of normal rainfall. If there is no rain for 48 hours after application, apply a second application. Do not use more than 200 lbs. per acre.

Apply 100 to 200 lbs. per acre in a band on each side of the seed furrow at planting. For established beans, apply 100 to 200 lbs. per acre in a band on each side of the seed furrow at planting.

side of the seed furrow at planting and as a side-dressing after plants become established. Allow a minimum of 21 days between applications. Do not apply to plants that are stressed.

Wheat: For control of aphids, hoppers, rips, mites and rips. Apply 100 to 200 lbs. (1 to 2 lbs. actual DI-SYSTON) per acre. For early season applications, apply 100 to 200 lbs. per acre in a band on each side of the seed furrow at planting. For established wheat, apply 100 to 200 lbs. per acre in a band on each side of the seed furrow at planting.

Call a crop consultant for more information. Do not use more than 200 lbs. per acre.

Highly toxic to fish and wildlife. Do not allow to drift to other crops. High concentration

Highly toxic to fish and wildlife. To protect water supplies, fish and wildlife, do not contaminate any body of water.

ATTN

Persons may result in ~~XXXXXXXXXX~~

May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Wash thoroughly after handling.

Symptoms of Poisoning: A sense of "tightness" in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhea.

Treatment: In case of poisoning, CALL A PHYSICIAN IMMEDIATELY. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a table-spoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If

ATROPINE IS ANTIDOTAL.

GUARANTEED ANALYSIS

Nitrogen	
Organic (from urea)	44.0%
Available Phosphoric Acid	0.0%
Water Soluble Potash	0.0%

NOTICE

The principal active insecticidal ingredient of this product is patented under U. S. Patent No. 2,759,410 and Canadian Patent No. 518,067 of Chemagro Corporation, Kansas City, Missouri and the purchase of this product includes a limited license under said patents to use or reformulate the product only in the manner provided by the recommended applications on the present label. The purchaser obligates itself to notify Chemagro Corporation of any proposed use or reformulation of such licensed product not provided by said recommended application, and to obtain Chemagro's approval in writing before making any such use or reformulation of the licensed product. Any use or reformulation intended for use, or the licensed product in the home or garden field is specifically prohibited by said license.

Non-warranty: Seller makes no warranty of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risks of use or handling whether in accordance with directions or not.