

AMERICAN CHEMICAL COMPANY
SALT LAKE CITY, UTAH


WALCO CAPTAN 7-1 DUST
(An Agricultural Fungicide)

ACTIVE INGREDIENTS:

Captan (N-trichloromethyl-4-chlorobenzene-1,3-dicarboximide) 7.5%

INERT INGREDIENTS:..... 92.5%
Total 100.0%

U. S. Pat. Nos. 2,553,770-2,553,771-2,553,776



Not licensed for use or export outside of the U. S., or its territories and possessions, Canada or Mexico.

U.S.D.A. Reg. No. 168-

Net Contents:

CAUTION: Keep out of reach of children.
Avoid inhaling dust. Avoid contact with skin.
Avoid contamination of feed and foodstuffs.
Observe the following use precautions.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. Observe the limitations given on label for specified crops.

Consult State Agricultural Experiment Station or State Extension Service authorities for additional information, as the timing, number and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

For best results, dust when air is calm. A good hand or power dusting machine should be used to secure completion of coverage of plant with a light film of dust. Begin application prior to or at first sign of infection. Repeat applications as necessary. Keep new growth protected.

RECOMMENDATIONS

APPLE: Scab, Lungeworm leaf spot-- to 50 lbs. per acre as required to maintain adequate coverage in the pre-bloom and blossom applications.

CHERRY: Brown Rot, Leaf Spots--25 to 50 lbs. per acre in the blossom and pre-harvest applications.

PEACH: Brown Rot and Scab-- to 50 lbs. per acre in the blossom and pre-harvest applications.

STRAWBERRIES: Botrytis-- to 50 lbs. per acre beginning as soon as new growth starts. Repeat at about weekly intervals, up to harvest. Apply after each picking if conditions favor fruit rot.

BEANS, PEAS: Soil Treatment--Damp off (Beans only), Root Rot--Work into top 3 or 4 inches of soil before planting. Apply 70 lbs. per acre.

WALCO CHEMICAL COMPANY
SALT LAKE CITY, UTAH


WALCO CAPLAN 7.5 DUST
(An Agricultural Fungicide)

ACTIVE INGREDIENTS:
Captan (N-trichloromethyl-2,2-dimethyl-1,3-dioxane-4,7-dicarboximide)..... 7.5%

INERT INGREDIENTS:..... 92.5%

Total 100.0%

U. S. Pat. Nos. 2,553,770-2,553,771-2,553,776



Not licensed for use or export outside of the U. S., or its territories and possessions, Canada or Mexico.

U.S.D.A. Reg. No. 168-

Net Contents:

CAUTION: Keep out of reach of children.
Avoid inhaling dust. Avoid contact with skin.
Avoid contamination of feed and foodstuffs.
Observe the following use precautions.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. Observe use limitations given on label for specified crops.

Consult State Agricultural Experiment Station or State Extension Service authorities for additional information, as the timing, number and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

For best results, dust when air is calm. A good hand or power dusting machine should be used to secure completion of coverage of plant with a light film of dust. Begin application prior to or at first sign of infection. Repeat applications as necessary. Keep new growth protected.

RECOMMENDATIONS

APPLE: Scab, Ringeye leaf spot--25 to 50 lbs. per acre as required to maintain adequate coverage in the pre-bloom and blossom applications.

CHERRY: Brown Rot, Leaf Spot--25 to 50 lbs. per acre in the blossom and pre-harvest applications.

PEACH: Brown Rot and Scab--25 to 50 lbs. per acre in the blossom and pre-harvest applications.

STRAWBERRIES: Botrytis--25 to 50 lbs. per acre beginning as soon as new growth starts. Repeat at about weekly intervals, up to harvest. Apply after each picking if conditions favor fruit rot.

BEANS, PEAS: Soil Treatment--Dorm off (Beans only), Root Rot--Work into top 3 or 4 inches of soil before planting. Apply 70 lbs. per acre.

AGRI-CHEMICAL COMPANY
ATLANTA, GEORGIA

AGRI-CHEMICAL COMPANY
AGRI-CHEMICALS DIVISION

PEARS: Damp-off, Root Rot, Black Rot--Apply 2 to 3 lbs. per acre when all seeds have begun germination. Repeat if necessary.

CUCUMBERS: Anthracnose, Downy Mildew, Watermelon, Honeydew Melons, Squash: Angular Leaf Spot, Anthracnose, Downy Mildew. Start applications when runners begin to form or at the first sign of disease. Apply at rate of 20 to 30 lbs. per acre on young plants increasing to 30 to 40 lbs. per acre on mature plants.

CUCUMBERS: Early and Late Blight--Make first application when plants are 2 to 4 inches high. Repeat at 7 to 10 days intervals as long as disease threatens. Apply 30 to 35 lbs. per acre on young plants. Use 40 lbs. per acre on mature plants.

POTATO SEED PIECE TREATMENT: Damp-off, Seed Rot--Apply at rate of 1 1/2 lbs. per 100 lbs. of seed pieces. Thoroughly coat each potato seed piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed store treated seed pieces in open crates 2 or more days before bagging.

TOMATOES: Anthracnose, Early and Late Blight, Gray Leaf Spot, Septoria Leaf Spot--Apply 30 to 40 lbs. per acre at weekly intervals on young plants; 40 to 50 lbs. per acre on mature plants.

SOIL AND GREENHOUSE BENCH TREATMENT--2 1/2 lbs. per 1,000 sq. ft. (100 lbs. per acre.)

Work into upper 3 or 4 inches of soil before planting. For damp-off and root rot disease of seedlings and transplants of roses, other shrubs, flowers, lawn seedbeds and vegetables such as bean, pea, celery, cabbage, egg plant, pepper and tomato.

ORNAMENTALS: Rose, Black Spot, Carmine Rust, Leaf Spot, Chrysanthemum, Flower Blight--Apply thoroughly prior to or at first signs of disease. Repeat application at 7 to 10 day intervals. Shorten interval during periods of frequent rain and heavy dews.

LAWNS, TURF: Brown patch, Damp-off, Leaf Spot, Root Rot--Apply 2 to 3 lbs. per 1,000 sq. ft. Repeat at 7 to 10 day intervals as needed. Melting snow--Apply 2 to 3 lbs. per 1,000 sq. ft. beginning when grass resumes growth in the Spring (March, April). Repeat at 7 to 14 day intervals as long as cool, wet weather persists.

CAUTION: This spray is toxic to fish, birds, bees, cats, dogs, and other animals. Do not apply to water, swimming pools, or to animals.

NON-WARRANTY: The use of this product is warranted only when used as directed on the label. The use of this product is not warranted when used in a manner other than that intended. The use of this product is not warranted when used in a manner other than that intended. The use of this product is not warranted when used in a manner other than that intended.