

ACCEPTED

MAY 16 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 168-387

MORGRO

CAUTIONS

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and food-stuffs. DO NOT REUSE CONTAINER. Destroy it by burying with wastes or burning. Stay away from smoke and fumes.

KEEP OUT OF REACH OF CHILDREN

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NON-WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.

EPA Est. 168-UT-1

DIAZINON 5% GARDEN DUST

160171 A

**KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON LEFT SIDE PANEL**

ACTIVE INGREDIENT:
Diazinon (4-methyl-6-pyrimidinyl)
NET WEIGHT: 5.00%
9.00%
100.00%

*U.S. Patent No. 2,784,843 Reg. T.M. Diazinon of Geigy Company.

EPA Reg. No. 168-387

TO USE AS A DUSTER: Punch out lid liner. Hold container in horizontal position and squeeze container. Changing position of container varies amount discharged.

DIRECTIONS FOR BEST RESULTS

- 1. Read the CAUTIONS on usage of this product.
- 2. Take cap off and pour into duster when using the refill container.
- 3. Apply the dust to the underside, as well as top side of all plants treated, where pests first appear. Repeat if needed per directions and cautions.
- 4. Apply the dust until you see a light film appear on all surfaces of the treated plants.
- 5. After application, wash all exposed skin surfaces with soap and water.

SOIL APPLICATION: Prior to planting the following vegetables: Beans, Broccoli, Cabbage, Carrots, Cauliflower, Corn, Cucumbers, Endive, Lettuce, Melons, Onions, Parsnips, Peas, Peppers, Potatoes, Radishes, Spinach, Squash, Tomatoes.

TO CONTROL CUTWORMS, SEED-CORN MAGGOTS, WIREWORMS: Apply evenly as a broadcast treatment over the surface of the soil. The amount in this package (1 lb.) should be broadcast over 550 square feet. Immediately after spreading, mix thoroughly with the soil to a depth of 6 to 8 inches.

FOLIAGE APPLICATION: Vegetables, Fruit and Flowers - Apply a thin film of dust to upper and lower leaf surfaces. Figures indicated after each vegetable, fruit or berry indicate the number of days before harvest of edible parts that dusting must be stopped:

- Beans (7), Peas (day of harvest), Potatoes (35), Tomatoes (3), for aphids and leafhoppers
- Beets (14), Broccoli (5), Cabbage (7), Cauliflower (5), Carrots (10), Radishes (10), for aphids.
- Onions (10), Cucumbers (7), Summer Squash (7), Winter Squash (3), Melons (3), for aphids, thrips, leafminers, mites.
- Strawberries (5), for aphids, cyclamen and two-spotted mites.
- Roses, Flowers, Shrubs, Ornamentals, for aphids, mites, leafhoppers, mealy bugs, white fly, scale crawlers.

USE PRECAUTIONS

DO NOT USE ON certain ferns, poinsettias, hibiscus, and gardenias since injury may occur. Not effective against certain spider mites on evergreens.

MORGRO

ACCEPTED

MAY 16 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 168-387

CAUTIONS

- Harmful if swallowed or inhaled. Avoid breathing dust.
- Avoid contact with skin and eyes. Wash thoroughly after handling.
- Change contaminated clothing daily.
- Avoid contamination of feed and food-stuffs.
- DO NOT REUSE CONTAINER. Destroy it by burying with wastes or burning. Stay away from smoke and fumes.

KEEP OUT OF REACH OF CHILDREN

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NON-WARRANTY Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.

EPA Est. 168-UT-1

DIAZINON 5% GARDEN DUST

160171 A

**A MULTIPURPOSE INSECT DUST FOR PESTS
IN SOIL AND ON FOLIAGE OF VEGETABLES,
FRUITS, FLOWERS AND SHRUBS**

CAUTION:

**KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON LEFT SIDE PANEL**

ACTIVE INGREDIENT:

*0,0-diethyl 0-(2-isopropyl -4-methyl-6-pyrimidinyl) phosphorothioate	5.00%
INERT INGREDIENTS:	95.00%
TOTAL	100.00%

*U.S. Patent No. 2,754,243 Reg. T.M. Diazinon of Geigy Company.

EPA Reg. No. 168-387

NET WEIGHT 1 LB.



TO USE AS A DUSTER: Punch out lid liner. Hold container in horizontal position and squeeze container. Changing position of container varies amount discharged.

**DIRECTIONS
FOR BEST RESULTS**

- 1 Read the CAUTIONS on usage of this product
- 2 Take cap off and pour into duster when using the refill container.
- 3 Apply the dust to the underside, as well as top side of all plants treated, where pests first appear. Repeat if needed per directions and cautions.
- 4 Apply the dust until you see a light film appear on all surfaces of the treated plants.
- 5 After application, wash all exposed skin surfaces with soap and water.

SOIL APPLICATION: Prior to planting the following vegetables: Beans, Broccoli, Cabbage, Carrots, Cauliflower, Corn, Cucumbers, Endive, Lettuce, Melons, Onions, Parsnips, Peas, Peppers, Potatoes, Radishes, Spinach, Squash, Tomatoes.

TO CONTROL CUTWORMS, SEED-CORN MAGGOTS, WIREWORMS: Apply evenly as a broadcast treatment over the surface of the soil. The amount in this package (1 lb.) should be broadcast over 550 square feet. Immediately after spreading, mix thoroughly with the soil to a depth of 6 to 8 inches.

FOLIAGE APPLICATION: Vegetables, Fruit and Flowers — Apply a thin film of dust to upper and lower leaf surfaces. Figures indicated after each vegetable, fruit or berry indicate the number of days before harvest of edible parts that dusting must be stopped.

- Beans (7), Peas (day of harvest), Potatoes (35), Tomatoes (3) for aphids and leafhopper.
- Beets (14), Broccoli (5), Cabbage (7), Cauliflower (5), Carrots (10), Radishes (10), for aphids.
- Onions (10), Cucumbers (7), Summer Squash (7), Winter Squash (3), Melons (3) for aphids, thrips, leafminers, mites.
- Strawberries (5), for aphids, cyclamen and two-spotted mites.
- Roses, Flowers, Shrubs, Ornamentals, for aphids, mites, leafhoppers, mealy bugs, white fly, scale crawlers.

USE PRECAUTIONS

DO NOT USE ON certain ferns, poinsettias, hibiscus, and gardenias since injury may occur. Not effective against certain spider mites on evergreens.