

WEEDS CONTROLLED

This product is a contact herbicide useful for controlling broadleaf weeds and grasses, such as puncturevine, wild mustard, wild oats, chickweed, sandbar, common ragweed, pigweed, lambsquarters, common plaintain, prostrate spurge, barnyardgrass, giant foxtail, yellow foxtail and to top-kill certain perennials such as Johnsongrass, Dal grass and nutsedge.

LAWN RENOVATION

The area to be treated should be mowed before treatment. Reapply five days later if green areas reappear. When foliage is all brown, matted areas should be raked and bare areas scarified. About five days after last treatment, water thoroughly and then seed at rate recommended for new lawns. Seed must be raked into soil. Keep seeded areas well watered.

CAUTION

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil. Do not track from treated to untreated areas. Do not contaminate waters used for domestic consumption, or by animals, wild-life, and aquatic life, or for irrigation purpose. Do not graze treated areas or feed treated forage to livestock.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

ANTIDOTE

If taken internally, induce vomiting and call physician at once.

Product No. 163705

GENERAL INFORMATION

Morgro Joint Grass & Weed Killer is recommended for the control of a wide variety of unwanted vegetation around buildings, along sidewalks, driveways and around trees and ornamentals. It is also recommended for complete lawn renovation. It is inactivated on contact with soil.

DIRECTIONS FOR USING

We suggest using a sprinkling can or similar applicator. Use this mixture of Joint Grass & Weed Killer as is without further dilution.

BE SURE AND SPRAY ONLY THE VEGETATION YOU WANT TO KILL. Wet the foliage to the point of run off. One quart will treat 100 sq. ft. of clipped or mowed weeds.

FOR GENERAL WEED CONTROL

Spray the unwanted vegetation thoroughly to the point of run off. Wilting will appear in 1 to 3 days and turn brown in 5 to 7 days. Repeat application may be necessary on the deep rooted hard to kill perennials. Best kill is obtained when the plants are young and during the hot sunny weather.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically recommended on this label.

\$ 3.98



JOINT GRASS-WEED KILLER

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON RIGHT SIDE PANEL**

ACTIVE INGREDIENTS:

Sodium Cacodylate 1.98%
Dimethylarsenic Acid (Cacodylic Acid) 0.34%

INERT INGREDIENTS 97.68%

TOTAL 100.00%

Total arsenic (as elemental) all in water soluble form 1.11%
Dimethylarsenic Acid equivalent per gallon 0.190 lbs. or 2.05%

USDA REG. NO. 168-404

NET CONTENTS: ONE GALLON

READY TO USE
NO MIXING - NON-SELECTIVE

READ ENTIRE LABEL. USE STRICTLY IN
ACCORDANCE WITH LABEL DIRECTIONS
AND CAUTIONS.

