WEEDS CONTROLLED

This product is a contact herbicide useful for controlling broadleaf weed. and grasses, such as puncturevine, wild mustard, wild oats, chickweed, sandbur, common ragweed, pigweed, lambsquarter, common plaintain, prostrate spurge, barnyardgrass, giant foxtail, yellow foxtail and to top-kill certain perennials such as Johnsongrass, Dallisgrass and nutsedge.

LAWN RENOVATION

The area to be treated should be mowed before treatment. Reapply five days later if green areas reappear. When foliage is all brown, matted areas should be raked and bare areas scarified. About five days after last treatment, water thoroughly and then seed at rate recommended for new lawns. Seed must be raked into soil. Keep seeded areas well watered.

CAUTION

Harmful if swallowed Avoid contact with skin. Avoid breathing spray mist Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil. Do not track from treated to untreated areas. Do not contaminate waters used for domestic consumption or by animals wild life, and aquatic life or for irrigation purpose.

Do not graze treated areas or feed treated forage to livestock.

CONTAINER DISPOSAL Do not reuse empty container Wash thoroughly with water and detergent crush if possible and discard in a safe place.

ANTIDOTE

If taken internally induce vomiting and call physician at once.

Product No. 163705

GENERAL INFORMATION

Morgro Joint Gress & Weed Killer is recommended for the control of a wide variety of unwented vegetation around buildings, along sidewalks, driveways and around trees and ornamentals. It is also recommended for complete lawn renovation. It is inactivated on contact with soil

DIRECTIONS FOR USING

We suggest using a sprinkling can or similar applicator. Use this mixture of Joint Grass & Weed Killer as is without further dilution.

BE SURE AND 'PRAY ONLY THE VEGETATION YOU WANT TO KILL. Wet the foliage to the point of run off. One quart will treat 100 sq. ft. of clipped or mowed weeds.

FOR GENERAL WEED CONTROL

Spray the unwaited vegetation thoroughily to the poin of run off. Wilting will appear in 1 to 3 days and turn brown in 5 to 7 days. Repeat application may be necessary on the deep rooted hard to kill perennials. Best kill is obtained when the plants are young and during the hot sunny veather.

NON-WARRANTY Manufacturer or seller is not liable for any injury or damage caused by the product due to misuse mishandling or any application not specifically recorded on this label.

\$ 3.98



READY TO USE NO MIXING - NON-SELECTIVE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

JOINT GRASS-WEED KILLER

CAUTION: KEEP OUT OF REACH OF CHILDREN SEE OTHER CAUTIONS ON RIGHT SIDE PANEL

Dimethylarsenic Acid equivalent per gallon 0 190 lbs. or 2.05% USDA REG NO 168-404 NET CONTENTS: ONE GALLON

WASATCH CHEMICAL COMPANY

