ACCEPTED

MAR 28 1969

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTRATION ED UNDER EPA REGISTRATION TO ATTACHED CONDITIONS.

WASCO

6% DITHANE M-45 DUST

ACTIVE INGREDIENT:

*A coordinated product of zinc ion and manganese ethylene bisdithiocarbamate....	4.8%
INERT INGREDIENT	95.2%
Total	100.0%

*From Dithane M-45 - Reg. T.M. Rohm and Haas Co.

USDA Reg. No. 168-

CAUTION: KEEP OUT OF REACH OF CHILDREN
May cause irritation to nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact wash with water; for eyes, wash copiously with water and get medical attention.

Contamination of fish-bearing waters with this product may harm fish.

IMPORTANT: Store in a cool, dry place. Do not allow to become wet or overheated in storage. Keep containers closed when not in use.

WASCO

6% DITHANE M-45 DUST

DIRECTIONS FOR USE

WASCO 6% DITHANE M-45 DUST can be used for control of many common diseases of vegetables and field crops. Application should be[illegible] when plants are small and when disease first appears. Rates [illegible] increase as plants become larger.

POTATO SEED-PIECE - For control of Fusarium seedpiece decay, [illegible] borne common scab and blackleg and to aid healing and suberization of cut seed surfaces: Use 1 lb. of WASCO 6% DITHANE M-45 DUST per 100 lbs. of cut or whole potato seed pieces. Apply the dust [illegible] manner that the entire surface of the seed piece becomes [illegible] covered with the fungicide dust. Plant [illegible] as soon after treatment as possible. Do not use treated seed potatoes [illegible] feed purposes.

CUCUMBERS, WATERMELONS, CANTALOUPES, [illegible], MUSKMELONS, [illegible] SQUASH - Downy mildew, anthracnose, scab, Alternaria leaf [illegible] Cercospora leaf spot and gummy stem blight: Use WASCO 6% DITHANE M-45 DUST at 25-45 lbs./acre. Begin applications when plants emerge or when disease is noted or expected in the area [illegible] at 5 to 7 day intervals.