

CAUTION

KEEP OUT OF REACH OF CHILDREN AT ALL TIMES
STORE IN A DRY PLACE.
DO NOT APPLY WHEN FRUIT TREE HAS FRUIT SET OR HANGING.

SMALL TREES MAY BE INJURED IF THE SOIL IS VERY DRY AND THE TREES ARE SHOWING DROUGHT STRESS.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label.

Manufactured by:
WASATCH CHEMICAL COMPANY
Subsidiary of Mountain Fuel Supply Company
Salt Lake City, Utah 84106



\$1.19

ROOT BORER CONTROL

A SOIL FUMIGANT FOR PEACH TREE BORER CONTROL ON PEACHES, CHERRIES, APRICOTS, PLUMS, NECTARINES
ACTIVE INTREDIENTS: Paradiichlorobenzene 99%
INERT INGREDIENTS: 1%
TOTAL 100%

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE LEFT PANEL FOR OTHER CAUTIONS.

USDA Reg No. 168-423 NET CONTENTS: 1 lb. (16 oz.)

FOR BEST RESULTS: Apply in the fall or early spring before fruit has set on tree. Clean away the trash, debris or soil around the base of the tree until the borer holes are exposed. Use 1 oz (2 level tablespoons) per tree for any of the tree varieties listed which will be bearing fruit. Apply the borer crystals evenly in a band around the tree base and at least 2" away from the trunk. Cover the crystals with soil in a cone shape and pack with the back of a spade to prevent the gas from escaping or being washed away. Normal watering, or having good moisture levels before applications, improves results.

ROOT BORER CONTROL penetrates the borer chambers in the form of gas which is toxic to peach tree borers. It is most effective when applied in the fall. This is the time when most borer eggs hatch and when the larvae are very young.

For younger trees, or in areas of extreme drought, consult with the State Agricultural Experiment Station or State Extension Service Specialists for special instructions to time amount to use and other requirements.

