

8 - 10 - 4 •

MORGRON[®]

the complete fertilizer

Aug 6, 1972

108 435 Comments

CRABGRASS & SPURGE KILLER

ACCEPTED WITH COMMENTS

CONTROLS WEED PESTS IN LAWNS AND ORNAMENTAL P

Cameraman's Note
Poor Copy

COVERS
2800 SQ. FT.
SEE DIRECTIONS
ON BACK

CONTAINS
DACTH



* DACTH is a
of Chemical Co.

**CAUTION. KEEP OUT OF REACH OF CHILDREN.
SEE BACK PANEL FOR REQUIRED CAUTIONS.**

AGENT WORKSHEET: *Handy/ color/ tubulose/ 100% cap* 1.27%
NET WEIGHT: 98.73%
TOTAL 100.00%

NEEDS CONTROLLED: *crabgrass* *Hairy Green foxtail*
CRABGRASS (Smooth and Hairy), GREEN FOXTAIL,
YELLOW FOXTAIL, LOWLAND, WICKERGRASS, FUELAGE, CANE-WEED, COMMON
CHICKEN, LAMPROUS, ANNUAL BURNING, BANNING GRASS, RED-ROOT FIG-
WEED, SPINY FICHEL, HISSING SPURGE and PROSTRATE SPURGE.

Cameraman's Note
Poor Copy

EPH 883. NO. 146-433

NET WEIGHT: 50 LBS.

MORGRO 6-10-4 CRABGRASS & SPURGE KILLER

GUARANTEED ANALYSIS

<p>TOTAL NITROGEN 6.0%</p> <p>1.5% Nitric Nitrogen Available From Ammonium Nitrate</p> <p>4.5% Ammonic Nitrogen From Ammonium Sulphate, Ammonium Nitrate, Ammonium Phosphate</p>	<p>AVAILABLE PHOSPHORIC ACID 10.0%</p> <p>From Potassium Phosphate</p> <p>POTASH SOLUBLE 4.0%</p> <p>From Potassium Phosphate</p> <p>IRON (Expressed as Fe) 1.0%</p> <p>From 10% Iron Sulphate</p>
--	--

Cameraman's Note
Poor Copy

HOW TO USE

1. Pour MORGRO 6-10-4 CRABGRASS & SPURGE KILLER into regular fertilizer spreader. Set spreader and apply at rate per 1000 sq. ft.
2. Spread uniformly over lawn when grass is dry.
3. Wash into soil with thorough watering.

DIRECTIONS FOR USE

ESTABLISHED LAWNS: Broadcast at rate of 20 lbs. per 1000 sq. ft. Apply in late winter or early spring before crabgrass, spurge and other annual grasses and broadleaf weeds germinate. Apply again in late summer or early fall at rate of 27 lbs. per 1000 sq. ft. to supplement early spring application and to control annual bluegrass and other erratic or late germinating grasses and weeds. Consult Agricultural Experiment Station or Extension Service Weed Specialist for local directions. **BE SURE TO APPLY BEFORE WEED SEEDS GERMINATE. WATER AFTER APPLICATION.**

NEWLY PLANTED LAWNS

Apply at 20 lbs. per 1000 sq. ft. AFTER NEW LAWN HAS BEEN MOWED AT LEAST TWICE.

SPREADER SETTINGS

MODEL	SETTING	MODEL	SETTING
Jackson #26-56	7	Whirlcast	8
Lawn Beauty #18-24-30-36	7	Scotts #35-75	4
Morcro	7	Sears Push Wheel Type	7

CHECK YOUR SPREADER: Is it feeding uniformly? Make sure it opens and closes properly. Wash out after using and hang in a dry place. Record proper setting for use next time.

Apply product when foliage of grass is dry. Water normally after treatment.

KEEP CHILDREN AND ANIMALS OFF treated areas until material has washed into soil and grass is completely dry.

CAUTIONS

HARMFUL IF SWALLOWED. WASH AFTER HANDLING OR USING.

AVOID INHALATION OF DUST AND CONTACT WITH EYES.

DO NOT CONTAMINATE FEED OR FOODSTUFFS.

TOXIC TO BIRDS AND FISH. USE WITH CARE.

DO NOT APPLY TO DICHOONDRA OR BENTGRASS.

DO NOT REUSE CONTAINER. DESTROY BY BURYING WITH WASTE OR BURNING IN A SAFE PLACE. STAY AWAY FROM SMOKE OR FUMES.

USE IN ORNAMENTALS

MORGRO 6-10-4 CRABGRASS & SPURGE KILLER provides effective pre-emergence control of crabgrass, annual grasses and certain broadleaf weeds in these ornamentals: Azalea, Barberry, Euonymus, Forsythia, Japanese Andromeda, Juniper, Lantana, Pachistima, Pine Rose, Tree Peony, Variegated Privet, Wilson Ivy, Yew, Ageratum, Alyssum, Aster, baby's breath, Bleeding Heart, Chrysanthemum, Candle Larkspur, Coleus, Columbine, Coral Bells, Dahlia, Evening Primrose, Fernleaf Yarrow, Feverfew, Forget-Me-Not, Gladiolus, Golden Marguerite, Golden Tuft, Lupine, Marigold, Mourning Bride, Mourning Pinks, Petunia, Poker Plant, Purple Coneflower, Rose Moe, Salvia, Scarlet Sage, Snapdragon, Verbena, Zinnia.

HOW TO APPLY: A complete, uniform application on a smooth soil surface, free from clods and weed seedling is essential to full control. Broadcast half the product in one direction and the other half at right angles to the first gives more uniform and complete coverage than a single, one-direction application. Water-in after treatment and wash residue from plant foliage. DO NOT USE on soils to be planted with seed the same season.

Use as a pre-emergent herbicide under U.S. Patent No. 2,923,004.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mis-handling or any application not specifically recommended on this label.

Manufactured by

WASATCH CHEMICAL DIVISION

Ei → entrada Industries inc.

P.O. BOX 6219 • SALT LAKE CITY, UTAH 84106

Cameraman's Note
Poor Copy