

**PRE-HARVEST AND GRAZING
USE LIMITATIONS**

Allow 1 day between application and harvest of asparagus and strawberries.
Allow 3 days between application and harvest of broccoli, Brussel sprouts, cabbage, cauliflower, head lettuce, garden beets (roots), radishes.
Allow 7 days between application and harvest of blackberries, boysenberries, dewberries.
Allow 14 days between last application and harvest of sugar beets, Chinese cabbage, endive (escarole), garden beets, leaf lettuce, spinach, Swiss chard.

VEGETABLE INSECT CONTROL

Beans: Green beans, lima beans, snap beans, cowpeas and black-eyed peas. Use for Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar and western bean cutworm.

Corn: Use for corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

Cucumber, Melons, Pumpkin and Squash: Use for pickleworm and melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

Potato, Tomato, Eggplant and Pepper: Use for Colorado potato beetle, flea beetles and leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs.

Cabbage, Broccoli, Brussel Sprouts & Cauliflower: Use for flea beetles, harlequin bug and leafhoppers. To control six-spotted leafhopper (aster yellows vector) apply on a 3 to 7 day schedule. Use for armyworms, cabbage worms, corn earworm, tarnished plant bug and stink bugs.

Where cabbage looper is the principal problem, use an alternative pesticide that is specifically recommended. This product will control only small cabbage loopers.

Carrots, Chinese Cabbage, Endive (Escarole), Garden Beets, Lettuce, Radishes, Spinach & Swiss Chard: Use for flea beetles, harlequin bug and leafhoppers. To control six-spotted leafhopper (aster yellows vector) apply on a 5 to 7 day schedule.

Use for armyworms, corn earworm, tarnished plant bug and stink bugs. For optimum control of corn earworm after lettuce heads begin to form, use on a 5 to 7 day schedule. Where cabbage looper is the principal problem, use an alternative pesticide that is specifically recom-

89¢



SEVIN* 5% DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)..... 5%

INERT INGREDIENTS:..... 95%

100%

**CAUTION: KEEP OUT OF REACH
OF CHILDREN
SEE OTHER CAUTIONS
ON RIGHT PANEL**

U.S.D.A. Reg. No. 168 441

Net Contents: 1 pound

**READ ENTIRE LABEL. USE IN STRICT ACCORDANCE
WITH LABEL DIRECTIONS AND CAUTIONS**

TO USE AS DUSTER

Punch out lid liner. Hold container in horizontal position and squeeze container. Changing position of the container varies the amount discharged.

FOR BEST RESULTS:

1. Apply the dust to the underside, as well as the top, of all plants treated, where pests first appear. Repeat if needed per directions and cautions.
2. Apply the dust until a light film appears on surfaces of the treated plants.
3. Read the caution on usage of this product.

NOTE

1. Some phytotoxicity may occur on some crops in the presence of rain or high humidity at several days following application.
2. Does not control spider mites, but is compatible with all common miticides.
3. This product is highly toxic to bees, exposed to direct treatment or residues on crops. Protect a information may be obtained from your Cooperative Agricultural Extension Service.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do Not Breathe Dust. Do Not Take Internally. Avoid Breathing of Dust. Skin Contact May Be Harmful. Avoid Contact. Wash Hands Before Eating. Take Shower or Bath after Work. Wear Regular Long-Sleeved Work Clothing. Change to Clean Clothing Daily.

NOTE TO PHYSICIAN: Carbaryl is a carbamate, reversible cholinesterase inhibitor. Atropine is antidote.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

*Trademark of Union Carbide Company for 1-naphthyl N-methylcarbamate.

NOTE

1. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days following application.
2. Does not control spider mites but is compatible with all common miticides.
3. This product is highly toxic to bees, especially to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CAUTION

HARMFUL IF SWALLOWED OR INHALED Do Not Breathe Dust. Do Not Take Internally. Avoid Breathing of Dust. Skin Contact May Be Harmful. Avoid Contact. Wash Hands before Eating. Take Shower or Bath after Work. Wear Respirator and Sieved Work Clothing. Change to Clean Clothing Daily.

NOTE TO PHYSICIAN: Carbyl is a moderately reversible cholinesterase inhibitor. Atropine is antidote.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

*Trademark of Union Carbide Company for
1-naphthyl N-methylcarbamate
U.S. Pat. No. 2,903,478
Product No. 167224

168-441

of infestation. Timing and good coverage are absolutely essential for effective control. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

Cucumber, Melons, Pumpkin and Squash: Use for pickleworm and melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

Potato, Tomato, Eggplant and Pepper: Use for Colorado potato beetle, flea beetles and leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bug and stink bugs.

Cabbage, Broccoli, Brussel Sprouts & Cauliflower: Use for flea beetles, harlequin bug and leafhoppers. To control six spotted leafhopper (aster yellows vector) apply on a 3 to 7 day schedule. Use for armyworms, cabbage worms, corn earworm, tarnished plant bug and stink bugs.

Where cabbage looper is the principal problem use an alternative pesticide that is specifically recommended. This product will control only small cabbage loopers.

Carrots, Chinese Cabbage, Endive (Escarole), Garden Beets, Lettuce, Radishes, Spinach & Swiss Chard: Use for flea beetles, harlequin bug and leafhoppers. To control six spotted leafhopper (aster yellows vector) apply on a 5 to 7 day schedule. Use for armyworms, corn earworm, tarnished plant bug and stink bugs. For optimum control of corn earworm after lettuce heads begin to form use on a 5 to 7 day schedule. Where cabbage looper is the principal problem use an alternative pesticide that is specifically recommended. For control of small cabbage loopers in home gardens, use on a 5 to 7 day schedule.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label.