

ACCEPTED  
 NOV 16, 1971  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR FERTILIZERS AND POISON REGISTER-  
 ED UNDER NO. 168-458 & SUBJECT  
 TO ALL OTHER COMMENTS.

## WARNING

KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or human beings. Do not permit children or pets to go onto the sprayed grass until spray has completely dried. Utensils such as teaspoons and table-spoons should not be used for food purposes after use with pesticides. KEEP OUT OF REACH OF DOMESTIC ANIMALS. DO NOT USE. POUR, SPILL OR STORE NEAR HEAT OR FLAME. Do not contaminate ornamental fishponds.

Product toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM TREATED AREAS. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply product ONLY as specified on this label.

SEE CHART for complete Insect List. Use dosages shown unless otherwise indicated:

## MEASURE INFORMATION

2 tablespoons = 1 fl. oz. or 6 teaspoons  
 1 cup = 8 fl. oz. or 1/2 pint.

2 tablespoons in 3 gals. water = 2 teaspoons in 1 gal. water or 1/2 pint in 24 gals. water.

(NOTE: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, and orchids since injury may occur.)

## WARNING: COMBUSTIBLE MIXTURE

## HOW TO MIX AND APPLY:

STEP 1: Determine size of spraying job. For lawns and vegetables measure area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear or before expected. Repeat treatments as needed or as noted in rate chart.

STEP 2: Use clean sprayer. Carefully measure and mix as recommended in chart.

STEP 3: Spray as directed in rate chart. Use coarse, drenching sprays for lawns, soil and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label.

\$6.98



DO NOT USE  
IN THE HOME

# DIAZINON LIQUID

(DIAZINON 25% EMULSIBLE)

CONTROLS LAWN AND GARDEN INSECTS

**WARNING: KEEP OUT OF REACH OF CHILDREN  
SEE SIDE PANELS FOR ADDITIONAL WARNINGS.**

ACTIVE INGREDIENTS:  
 O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl)  
 phosphorothioate ..... 25%  
 Aromatic Petroleum Derivative Solvent ..... 57  
 INERT INGREDIENTS: ..... 18  
 TOTAL ..... 100%

U.S. Patent  
No. 2,754,243

Spray Concentrate contains Diazinon\* Insecticide.  
 DIAZINON\* is a registered trademark of Geigy Chemical Corporation.

EPA Reg. No. 168-458

NET CONTENTS: ONE QUART (32 FL. OZ.)

MANUFACTURED BY  
**WASATCH CHEMICAL COMPANY**

A SUBSIDIARY OF MOUNTAIN FUEL SUPPLY CO.  
SALT LAKE CITY, UTAH 84106

## DIAZINON 25% EMULSIBLE INSECT CONTROL

Kills certain insects which destroy lawns, roses, flowers, trees and shrubs or that annoy pets and people. Contains DIAZINON\* insecticide with provides positive control of many "hard-to-kill" sucking and chewing insects. DIAZINON 25% EMULSIBLE works fast. Remains effective for an extended period of time.

CHART FOR APPLICATION OF DIAZINON 25% EMULSIBLE

Problem Area	Pests	Rate & Area Covered	Notes
Lawns: Bluegrass, Marion, Fescue, St. August- ine, Dichondra, Bent, Bermuda	Chinch Bugs*, Sod Webworms, Bermuda stem, Ants, Ticks, Earwig, Lawn Pillbugs	2 cups (1 fl. oz.) in 3 gals. of water covers 125 sq. ft. 8 fl. oz. in 24 gals. of water covers 1,000 sq. ft.	Thoroughly wet down grass a few hours before applying. For Lawn Pillbugs, water soak thoroughly after treat- ment. *In the north, if Chinch Bugs are the only insect problem, a lower rate of application can be used: 2 cups in 3 gals. of water covers 250 sq. ft.
Melons		4 cups (2 fl. oz.) in 3 gals. of water covers 125 sq. ft. 16 oz./24 gals. water covers 1,000 sq. ft.	
Rhodes-grass Scale Crawlers		2 cups (1 fl. oz.) in 3 gals. of water covers 100 sq. ft. 10 oz./30 gals. water covers 1,000 sq. ft.	Apply when Scale Crawlers first emerge
Roses and Flowers Roses, Azaleas, Chrysanthemums, Gladiolus, Carna- tions	Aphids, Mites, Leafhoppers, Thrips	2 cups (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray	Spray underside of leaves wet plants to the drooping point
Trees and Shrubs: Oak, Maple, Locust, Holly, Boxwood, Pine, Spruce, Willow	Aphids, Mites, Scale Crawlers, Leaf Miners, Borerworms	2 cups (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray.	Wet leaves and branches to the drooping point. Try to penetrate dense foliage
Fruits: Apples, Peaches, Oranges, Plums, Grapes, Citrus, Strawberries	Codling Moths, Mites, Aphids, Leaf Rollers, Scale Crawlers	2 cups (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Start spraying after bloom- ing and repeat at 10 to 14 day intervals if needed. Do not spray fruit within 21 days of picking
Vegetables: Tomatoes, Carrots, Potatoes, Beans, Cucumbers, Onions, Cantaloupes, Peppers, Cabbage	Mexican Bean Beetles, Aphids, Leaf Miners, Imported Celo beetworms, Cucumber Beetles	2 cups (1 fl. oz.) in 3 gals. of water covers 1,000 sq. ft. For carrots, cucumbers, cabbage & onions, use 1 1/2 cups in 3 gal- ons of water per 1,000 sq. ft. For peppers, reduce dosage to 1/2 cup in 3 gal. of water per 1,000 sq. ft.	Generally applications at 7 to 10 day intervals may be required. Do not apply less than one day of picking tomatoes. 30 days of picking potatoes, or 14 days of picking all other crops listed in this section.
Cabbage, Carrots, Coffin-leafers, Peas, Pole beans, Radishes, Red beans, Snap beans, Tom- atoes, Turnips	Cutworms (Sur- face and Sub- terranean), Wireworms, Root Maggots, Garden Syrphids, Mole Crickets	Apply 1 1/2 fl. oz. per 1,000 sq. ft. as a soil treatment in sufficient water to obtain thorough coverage.	Make application to soil surface as a broadcast spray prior to planting. Thoroughly mix the chemical into the top 4-6 inches of soil
Nuisance Pests in Outside Areas:	Flies, Ants, Cockroaches, Beetles, Fleas, Dove Mites, Spiders, Ticks, Chiggers, Earwigs	Add 8 fl. oz. per 3 gals. of water.	Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling four corners, walls, garbage cans, and wire cracks and other places where in- sects may hide. Do not treat plants or animals with this dosage rate

PRODUCT NO. 161308

168-458

11/16/1971

111