

**WARNING**

KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or human beings. Do not permit children or pets to go onto the sprayed grass until spray has completely dried. Utensils such as teaspoons and table-spoons should not be used for food purposes after use with pesticides. KEEP OUT OF REACH OF DOMESTIC ANIMALS. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR FLAME. Do not contaminate ornamental fishponds.

Product toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM TREATED AREAS. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply product ONLY as specified on this label.

SEE CHART for complete Insect List. Use dosages shown unless otherwise indicated:

**MEASURE INFORMATION**

2 tablespoons = 1 fl. oz. or 6 teaspoons  
 1 cup = 8 fl. oz. or 1/2 pint.  
 2 tablespoons in 3 gals. water = 2 teaspoons in 1 gal. water or 1/2 pint in 24 gals. water.  
 (NOTE: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, and orchids since injury may occur.)

**WARNING: COMBUSTIBLE MIXTURE**

**HOW TO MIX AND APPLY:**

STEP 1: Determine size of spraying job. For lawns and vegetables measure area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear or before expected. Repeat treatments as needed or as noted in rate chart.

STEP 2: Use clean sprayer. Carefully measure and mix as recommended in chart.

STEP 3: Spray as directed in rate chart. Use coarse, drenching sprays for lawns, soil and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

NON-WARRANTY. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label.

**\$6.98**



**DO NOT USE  
IN THE HOME**

**DIAZINON LIQUID**

(DIAZINON 25% EMULSIBLE)

**CONTROLS LAWN AND GARDEN INSECTS**

**WARNING: KEEP OUT OF REACH OF CHILDREN  
SEE SIDE PANELS FOR ADDITIONAL WARNINGS.**

<b>ACTIVE INGREDIENTS:</b>		
O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate	.....	25%
Aromatic Petroleum Derivative Solvent	.....	57
<b>INERT INGREDIENTS:</b>	.....	18
<b>TOTAL</b>	.....	<b>100%</b>

U.S. Patent No. 2,754,243  
 Spray Concentrate contains Diazinon\* Insecticide.  
 DIAZINON\* is a registered trademark of Geigy Chemical Corporation.

EPA Reg. No. 168-458

**NET CONTENTS: ONE QUART (32 FL. OZ.)**

MANUFACTURED BY  
**WASATCH CHEMICAL COMPANY**  
 A SUBSIDIARY OF MOUNTAIN FUEL SUPPLY CO.  
 SALT LAKE CITY, UTAH 84106

**DIAZINON 25% EMULSIBLE INSECT CONTROL**

Kills certain insects which destroy lawns, roses, flowers, trees and shrubs or that annoy pets and people. Contains DIAZINON\* insecticide with provides positive control of many "hard-to-kill" sucking and chewing insects. DIAZINON 25% EMULSIBLE works fast. Remains effective for an extended period of time.

**CHART FOR APPLICATION OF DIAZINON 25% EMULSIBLE**

Problem Area	Pests	Rate & Area Covered	Notes
Lawn Bluegrass, Mennon Fescue, St. August ine, Dichondra, Bent, Bermuda	Chinch Bugs, <sup>1</sup> Sod Webworms, Bermuda Weevils, Ants, Ticks, Emerald Lawn Pillbugs	2 tpsps (1 fl. oz.) in 3 gals. of water covers 125 sq. ft. 8 fl. oz. in 24 gals. of water covers 1,000 sq. ft.	Thoroughly wet down grass a few hours before applying for Lawn Pillbugs, water soak thoroughly after treat- ment. <sup>1</sup> In the north, if Chinch Bugs are the only insect problem, a lower rate of application can be used. 2 tpsps in 3 gals. of water covers 250 sq. ft.
	Meliponids	4 tpsps (2 fl. oz.) in 3 gals. of water covers 125 sq. ft. 16 oz. 24 gals. water cover 1,000 sq. ft.	
	Rhodes-grass Scale Crawlers	2 tpsps (1 fl. oz.) in 3 gals. of water covers 100 sq. ft. 10 oz. 30 gals. water covers 1,000 sq. ft.	Apply when Scale Crawlers first emerge.
Roses and Flowers Roses, Apples, Chrysanthemums, Gladiol, Carna- tions	Aphids, Meas. Leafhoppers, Thrips	2 tpsps (1 fl. oz.) per 3 gals. of water. Apply as a thorough leafy spray	Spray underside of leaves near points to the drooping point
Trees and Shrubs Oak, Maple, Locust, Holly, Beechwood, Pine, Spruce, Willow	Aphids, Meas., Scale Crawlers, Leaf Miners, Bagworms	2 tpsps (1 fl. oz.) per 3 gals. of water. Apply as a thorough leafy spray	Wet leaves and branches to the drooping point. Try to penetrate dense foliage
Fruit Apples, Peaches, Oranges, Plums, Grapes, Cherries, Strawberries	Codling Moths, Meas., Aphids, Leaf Rollers, Leaf Miners, Scale Crawlers	2 tpsps (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Start spraying after bloom- ing and repeat at 10 to 14 day intervals if needed. Do not spray near within 21 days of picking.
Vegetables Tomatoes, Carrots, Potatoes, Beans, Cucumbers, Onions, Cauliflower, Peppers, Cabbage	American Bean Beetles, Aphids, Leaf Miners, Imported Cab- bageworms, Cucumber Beetles	2 tpsps (1 fl. oz.) in 3 gals. of water covers 1,000 sq. ft. For carrots, cucumbers, cabbage & onions, use 1 1/2 tpsps in 3 gal. of water per 1,000 sq. ft. For peppers, reduce dosage to 1/2 tpsps in 3 gal. of water per 1,000 sq. ft.	Generally applications at 7 to 10 day intervals may be required. Do not apply with- in one day of picking tomatoes, 30 days of harvesting potatoes, or 14 days of picking all other crops listed in this section.
Cabbage, Carrots, Corn, Lettuce, Peanut Beans, Radishes, Red beets, Snap beans, Toma- tocs, Turnips	Cutworms (Sur- face and Sub- terranean), Wireworms, Root Maggots, Carrots Sympterygus, Mole Crickets	Apply 14 fl. oz. per surface in a broadcast spray prior to planting. Thoroughly mix the chemical into the top 4 inches of soil.	Make application to soil prior to planting. Thoroughly mix the chemical into the top 4 inches of soil.
Nuisance Pests in Outside Areas	Flies, Ants, Cockroaches, Blattelder Bugs, Clove Moth, Spiders, Ticks, Chiggers, Earwig	Add 8 fl. oz. per 3 gals. of water	Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling, foundations, walls, garbage cans, and into cracks and other places where in- sects may hide. Do not treat plants or animals with this dosage rate.

PRODUCT NO. 161308

ACCEPTED  
 AUG 3, 1972  
 UNDER THE FEDERAL INSECTICIDE  
 ACT AND SUBSIDIARY ACT  
 FOR ECONOMIC PEST CONTROL  
 REGISTERED UNDER NO. 168-458 SUBJECT  
 TO ALL OTHER COMMENTS.

168-458

08/03/1972

1/1