

# WARNING

KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or human beings. Do not permit children or pets to go onto the sprayed grass until spray has completely dried. Utensils such as teaspoons and table-spoons should not be used for food purposes after use with pesticides. KEEP OUT OF REACH OF DOMESTIC ANIMALS. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR FLAME. Do not contaminate ornamental fishponds.

Product toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM TREATED AREAS. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply product ONLY as specified on this label.

SEE CHART for complete insect List. Use dosages shown unless otherwise indicated.

## MEASURE INFORMATION

2 tablespoons = 1 fl. oz. or 6 teaspoons  
 1 cup = 8 fl. oz. or 1/2 pint.  
 2 tablespoons in 3 gals. water = 2 teaspoons in 1 gal. water or 1/2 pint in 24 gals. water.  
 (NOTE: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, and orchids since injury may occur.)

## WARNING: COMBUSTIBLE MIXTURE

### HOW TO MIX AND APPLY:

- STEP 1:** Determine size of spraying job. For lawns and vegetables measure area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear or before expected. Repeat treatments as needed or as noted in rate chart.
- STEP 2:** Use clean sprayer. Carefully measure and mix as recommended in chart.
- STEP 3:** Spray as directed in rate chart. Use coarse, drenching sprays for lawns, soil and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

DO NOT REUSE CONTAINER. Destroy when empty.

160182

EPA Est. 168-UT-1

ACCEPTED  
 168-458  
 JUN 11 1974  
 UNITED STATES DEPARTMENT OF AGRICULTURE  
 FUNGICIDE AND PESTICIDE DIVISION  
 FOR THE FEDERAL INSECTICIDE REGISTRATION ACT  
 EPA REG. NO. 168-458  
 SUBJECT: DIAZINON  
 TO ATTACHED COMMENTS.



DO NOT USE  
 IN THE HOME

# DIAZINON FRUIT & VEGETABLE SPRAY

(ALSO CONTROLS LAWN & GARDEN INSECTS)

**WARNING: KEEP OUT OF REACH OF CHILDREN  
 SEE LEFT PANEL FOR OTHER WARNINGS**

**ACTIVE INGREDIENTS:**

0.0-diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate	25.0%
Aromatic Petroleum Derivative Solvent	57.0%
<b>INERT INGREDIENTS:</b>	<u>18.0%</u>
<b>TOTAL</b>	<b>100.0%</b>

Spray Concentrate contains Diazinon\* Insecticide.  
 DIAZINON\* is a registered trademark of Geigy Chemical Corporation.

**EPA Reg. No 168-458      NET CONTENTS: ONE QUART (32 FL. Oz.)**



Kills certain  
 ants and per  
 many hard-t  
 work fast R

Prun Lawn By m Bent
Roses Roses Chrysi Gladio tions
Trees Oak, A Holly, Pine, S Willow
Fruits Apples Oranges Grapes Strawb
Vegeta Tomato Potato Cucum Cantal Pepper
Cabbag Corn, L Peas, F Radish Snap b Tues, T
Beans Couts

NON WARRANTY  
 use of this pro  
 is not liable for  
 application not

## WARNING

KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of food and foodstuffs. Do not use on household pets or human beings. Do not permit children or pets to go onto the sprayed grass until spray has completely dried. Utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides. KEEP OUT OF REACH OF DOMESTIC ANIMALS. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR FLAME. Do not contaminate ornamental fishponds.

Product toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM TREATED AREAS. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply product ONLY as specified on this label.

SEE CHART for complete Insect List. Use dosages shown unless otherwise indicated:

### MEASURE INFORMATION

2 tablespoons = 1 fl. oz. or 6 teaspoons

1 cup = 8 fl. oz. or 1/2 pint.

2 tablespoons in 3 gals. water = 2 teaspoons in 1 gal. water or 1/2 pint in 24 gals. water.

(NOTE: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, and orchids since injury may occur.)

### WARNING: COMBUSTIBLE MIXTURE

#### HOW TO MIX AND APPLY:

STEP 1: Determine size of spraying job. For lawns and vegetables measure area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear or before expected. Repeat treatments as needed or as noted in rate chart.

STEP 2: Use clean sprayer. Carefully measure and mix as recommended in chart.

STEP 3: Spray as directed in rate chart. Use coarse, drenching sprays for lawns, soil and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

DO NOT REUSE CONTAINER. Destroy when empty.

160182

EPA Est. 168-UT-1

ACCEPTED

168-458

JUL 11 1974

UNDER FEDERAL PESTICIDE  
FUNGICIDE AND RODENTICIDE  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. SUBJECT  
TO ATTACHED COMMENTS.

MURGRU

DO NOT USE  
IN THE HOME

# DIAZINON FRUIT & VEGETABLE SPRAY

(ALSO CONTROLS LAWN & GARDEN INSECTS)

Kills certain insects on  
pets and people  
many "hard-to-kill"  
works fast. R

NON-WARRANTY:  
use of this product  
is not liable for  
application not s

## DIAZINON FRUIT & VEGETABLE SPRAY

Kills certain insects which destroy lawns, roses, flowers, trees and shrubs or that annoy pets and people. Contains DIAZINON<sup>®</sup> insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. DIAZINON FRUIT & VEGETABLE SPRAY works fast. Remains effective for an extended period of time.

### CHART FOR APPLICATION

Problem Area	Pests	Rate & Area Covered	Notes
Lawns: Lawns, Roses, Flowers, Trees, Shrubs, etc.	Chinch Bugs, Billbugs, Grasshoppers, Crickets, Moles, Scale Insects	2 tbsps. (1 fl. oz.) in 3 gals. of water covers 125 sq. ft. 8 fl. oz. in 24 gals. of water covers 1,000 sq. ft.	Thoroughly wet down grass a few hours before spraying. For Lawn Billbugs, water lawn thoroughly after treat- ment. *In the north, if Chinch Bugs are the only insect problem, a lower rate of application can be used: 2 tbsps. in 3 gals. of water covers 250 sq. ft.
	Millipedes	4 tbsps. (2 fl. oz.) in 3 gals. of water covers 125 sq. ft. 16 oz./24 gals. water cover 1,000 sq. ft.	
	Worms—grass Scale Crawlers	2 tbsps. (1 fl. oz.) in 3 gals. of water covers 100 sq. ft. 10 oz./30 gals. water covers 1,000 sq. ft.	Apply when Scale Crawlers first emerge.
Roses and Flowers: Roses, Azaleas, Chrysantheums, Gladioli, Cam- panula	Aphids, Mites, Leafhoppers, Thrips	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray.	Spray underside of leaves. Wet plants to the dripping point.
Trees and Shrubs: Oak, Maple, Juniper, Hill, Silvered, Pine, Spruce, Willow	Aphids, Mites, Scale Crawlers, Leaf Miners, Bagworms	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray.	Wet leaves and branches to the dripping point. Try to penetrate dense foliage.
Fruits: Apples, Peaches, Grapes, Plums, Grapes, Cherries, Strawberries	Codling Moths, Mites, Aphids, Leaf Rollers, Scale Crawlers	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Start spraying after bloom- ing and repeat at 10 to 14 day intervals if needed. Do not spray fruit within 21 days of picking.
Vegetables: Tomatoes, Carrots, Potatoes, Beans, Cucumbers, Onions, Cauliflower, Peppers, Cabbage	Mexican Bean Beetles, Aphids, Leaf Miners, Imported Cab- bageworms, Cucumber Beetles	2 tbsps. (1 fl. oz.) in 3 gals. of water covers 1,000 sq. ft. For carrots, cucumbers, cabbage & onions, use 1 1/2 tbsps. in 3 gal. of water per 1,000 sq. ft. For peppers, reduce dosage to 1/2 tbsps. in 3 gal. of water per 1,000 sq. ft.	Generally applications at 7 to 10 day intervals may be required. Do not apply with- in one day of picking tomatoes, 35 days of harvesting potatoes, or 14 days of picking all other crops listed in this section.
Cabbage, Carrots, Corn, Lettuce, Pars, Pole beans, Radishes, Red Beets, Snap beans, Tom- atoes, Turnips	Cutworms (Sur- face and Sub- terranean) Wireworms Root Maggots Garden Symphylans, Mole Crickets	Apply 14 fl. oz. per 1,000 sq. ft. as a soil treatment in sufficient water to obtain thorough coverage.	Make application to soil surface as a broadcast spray prior to planting. Thoroughly mix the chemical into the top 4-6 inches of soil.
Nuisance Pests in Outside Areas <b>Vill 2</b>	Flies, Ants, Cockroaches, Blowflies, Bugs, Clover Mites, Spiders, Ticks, Chiggers, Earwigs	Add 8 fl. oz. per 3 gals. of water	Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling foundations, walks, garbage cans, and into cracks and other places where in- sects may hide. Do not treat plants or animals with this dosage rate.

**NON-WARRANTY:** Manufacturer or Seller makes no warranty express or implied concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.