

\$4.95

GENERAL INFORMATION

FORE* Turf Fungicide is a unique and different fungicide, which when used according to directions, provides very high fungicidal activity and is safe for use on turf grasses. This new fungicide provides proven control of most common turf grass diseases.

Diseases of turf grass can attack suddenly and unexpectedly. Severe damage may result, including total loss of large areas. A regular protective spraying program will minimize risk of disease damage and can be accomplished with lower rates and less frequent fungicide applications. If disease is established, high rates of fungicide and more frequent applications will be necessary for control. A regular protective program will provide maximum protection.

TO MIX...slowly sift FORE* Turf Fungicide into spray tank while it is being filled. Agitate.

CAUTIONS: KEEP OUT OF REACH OF CHILDREN

This product may cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact, wash with water. If contact with eyes, wash thoroughly with water. Get medical attention.

A. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.


B. Do not reuse container. Destroy when empty.

IMPORTANT

Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring about chemical changes which can impair the fungicide effectiveness of product and may also generate flammable vapors. Keep container closed when not in use.

NON-WARRANTY. Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any other application not specifically recommended on this label.

FORE* Registered Trade Mark, Rohm & Haas Company



FORE

TURF FUNGICIDE

TREATS UP TO 4,000 SQ. FT. OF LAWNS

CAUTION:

KEEP OUT OF REACH OF CHILDREN SEE OTHER CAUTIONS ON PANELS

ACTIVE INGREDIENTS

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate 80% in which the ingredients are:

Manganese + +	16%
Zinc + +	2%
Ethylene bisdithiocarbamate ion (C ₄ H ₆ N ₂ S ₄)	62%
INERT INGREDIENT	20%
TOTAL	100%

U.S. Patents 2,504,404; 2,974,156 and 3,379,610.

U.S.D.A. Reg. No. 168-462

NET WT: ONE POUND

DIRECTIONS FOR USE

TURF DISEASES: Fusarium Blight (F. roseum), Red Thread, Copper Spot, Stem Molds.

In a protective spray program use this product at 4 to 6 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in spring. Repeat every 7 to 14 days.

When conditions are especially favorable for disease development increase FORE Turf Fungicide rate to 6 to 8 ounces per 1000 sq. ft. Repeat applications every 7 days until disease threat passes.

TURF DISEASES: Helminthosporium Melting Out, Rhizoctonia Brown Patch, Rust.

Use FORE at 4 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in the spring and repeat every 7 to 14 days for Helminthosporium and Rust. Repeat every 7 days for Rhizoctonia con' d).

TURF DISEASE: Pythium Blight

Use FORE at 8 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications at first sign of disease and repeat at 5 day intervals, or oftener if conditions promote disease development. Continue until disease threat has passed.

TURF DISEASE: Algae Control in Turf Grass

Use FORE at 6 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when algae begin to appear. Repeat every 7 days until threat has passed.

TURF DISEASE: Dollar Spot (Sclerotinia)

In a protective spray program use FORE at 6 to 8 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens-up in spring and repeat every 7 to 14 days. When conditions are especially favorable for dollar spot development, use FORE at 8 ozs. per 1000 sq. ft. Repeat every 7 days until disease threat passes.

TURF DISEASE: Fusarium Snow Mold

Use FORE at 6 to 8 ozs. per 1000 sq. ft. in sufficient water to provide adequate coverage. Make applications during winter at 2 to 6 week intervals.

FOLLOW LOCAL TURF DISEASE RECOMMENDATIONS FOR FURTHER DETAILS ON APPLICATION TIMING

Product No. 164765

WASATCH CHEMICAL COMPANY

\$4.95

GENERAL INFORMATION

FORE® Turf Fungicide is a unique and different fungicide which when used according to directions provides very high fungicidal activity and is safe for use on turf grasses. This new fungicide provides proven control of most common turf grass diseases.

Diseases of turf grass can attack suddenly and unexpectedly. Severe damage may result including loss of large areas. A regular protective spraying program will minimize risk of disease damage and can be accomplished with lower rates and less frequent fungicide applications. If disease is established, high rates of fungicide and more frequent applications will be necessary for control. A regular protective program will provide maximum protection.

TO MIX: slowly sift FORE® Turf Fungicide into spray tank while it is being filled. Agitate.

CAUTIONS: KEEP OUT OF REACH OF CHILDREN

This product may cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact, wash with water. If contact with eyes, wash thoroughly with water. Get medical attention.

A. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.


B. Do not reuse container. Destroy when empty.

IMPORTANT

Keep away from fire and sparks. Store in a cool, dry place. Do not allow it to become wet or overheated in storage. This may bring about chemical changes which can impair the fungicide effectiveness of product and may also generate flammable vapors. Keep container closed when not in use.

NON WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any other application not specifically recommended on this label.

FORE® Registered Trade Mark, Rohm & Haas Company



FORE TURF FUNGICIDE

TREATS UP TO 4,000 SQ. FT. OF LAWN...

CAUTION: KEEP OUT OF REACH OF CHILDREN SEE OTHER CAUTIONS ON PANELS

ACTIVE INGREDIENTS

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate 80% in which the ingredients are:

Manganese + +	16%
Zinc + +	2%
Ethylene bisdithiocarbamate ion (C ₂ H ₆ N ₂ S ₄)	62%
INERT INGREDIENT	20%
TOTAL	100%

U.S. Patents 2,504,404; 2,974,156 and 3,379,610.

U.S.D.A. Reg. No. 168-462

NET WT: ONE POUND

MANUFACTURED BY WASATCH CHEMICAL COMPANY A SUBSIDIARY OF MOUNTAIN FUEL SUPPLY CO. SALT LAKE CITY, UTAH 84106

DIRECTIONS FOR USE

TURF DISEASES: Fusarium Blight, Fairy Ring, Red Thread, Copper Spot, Sclerotinia Molds

In a protective spray program use this product at 4 to 6 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens up in spring. Repeat every 7 to 14 days.

When conditions are especially favorable for disease development, increase FORE Turf Fungicide rate to 6 to 8 ounces per 1000 sq. ft. Repeat applications every 7 days until disease threat passes.

TURF DISEASES: Helminthosporium Melting Out, Rhizoctonia Brown Patch, Rust

Use FORE at 4 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens up in the spring and repeat every 7 to 14 days for Helminthosporium and Rust. Repeat every 7 days for Rhizoctonia control.

TURF DISEASE: Pythium Blight

Use FORE at 8 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications at first sign of disease and repeat at 5 day intervals, or oftener if conditions promote disease development. Continue until disease threat has passed.

TURF DISEASE: Algae Control in Turf Grass

Use FORE at 6 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when algae begin to appear. Repeat every 7 days until threat has passed.

TURF DISEASE: Dollar Spot (Sclerotinia)

In a protective spray program use FORE at 6 to 8 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage. Start applications when grass greens up in spring and repeat every 7 to 14 days. When conditions are especially favorable for dollar spot development, use FORE at 8 ozs. per 1000 sq. ft. Repeat every 7 days until disease threat passes.

TURF DISEASE: Fusarium Snow Mold

Use FORE at 6 to 8 ozs. per 1000 sq. ft. in sufficient water to provide adequate coverage. Make applications during winter at 2 to 4 week intervals.

FOLLOW LOCAL TURF DISEASE RECOMMENDATIONS FOR FURTHER DETAILS ON APPLICATION TIMING

Product No. 164765