## DIRECTIONS

Application should be made in Spring or Fall,2 to 4 weeks after emergence of legumes and when seedling weeds are less than 1 or 2 inches high or have rosettes less than 3 inches across. Later applications give unsatisfactory control.

Wet weeds thoroughly but not enough to cause runoff. Use 5 to 10 gallons of water per acre or sufficient for thorough coverage.

Seedling or Established Legumes such as alfalfa, red clover, trefoil and white clover:

Annual broadleaf weeds - cocklebur, kochia, lamb's quarter, nightshade, pigweed, ragweed, stinkweed (penny cress).

Less than I inch high - 1 to 2 qts.2,4-D Butyric/acre.

I to 3 inches high - 2 to 3 qts.2,4-D Butyric/acre.


Other broadleaf weeds - field bindweed (morning glory), seedling curled dock, Russian thistle, smartweed.

Up to 3 inches high - 2 to 3 qts.2,4-D Butyric/acre.



EPA Reg.No.168-474

WASATCH CHEMICAL DIVISION : ::
Ei→ entrada industries inic



## Established Legumes:

Other broadleaf weeds - Canada thistle, white top (hoary cress) and others listed above.

Up to 3 inches high - 3 qts. 2,4-D Butyric/acre.

CAUTION: Avoid contact with skin, eyes or clothing. Do not graze treated areas nor feed straw or hay from treated crops to livestock within 30 days after application. Avoid spray drift to susceptible plants. Coarse sprays are less likely to drift. Do not contaminate irrigation ditches or water used for domestic purposes. Do not use same spray equipment for other purposes, unless cleaned with a suitable chemical cleaner. Do not store near fertilizers, seed, insecticides or fungicides. Harmful if swallowed. Do not reuse container. Destroy by burying in a safe place. Do not burn.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label.

NET CONTENTS: 1 Gallon