

BENOMYL is highly effective against fungus diseases on roses, flowers, ornamentals and turf, providing both curative (eradicator) action and protective action. Mix with water to apply as spray. Addition of a surfactant to the spray mixture increases surface acceptance of fungicide on hard-to-wet plants such as roses.

ROSES (field & greenhouse): Powdery Mildew, Black Spot - use 1 Tbspn per 2 gals. water. Apply when disease first appears. Repeat at 10-14 day intervals during growing season. Shorten intervals during humid, rainy weather.

FLOWERS & ORNAMENTALS (field & greenhouse): Powdery Mildew, Botrytis Grey Mold - use 1 Tbspn per 2 gals. water. Apply when disease first appears. Repeat at 10-14 day intervals as long as needed.

TURF: Large Brown Patch, Dollar Spot - apply 2 oz. (12 Tbspns) per 1000 sq. ft. in sufficient water to cover (2.5 gals per 1000 sq. ft.). Apply when disease appears and continue at 10-14 day intervals. If conditions favor disease development, reduce interval to 5-7 days. DO NOT GRAZE OR FEED clippings from treated areas to livestock.

Distributed by WASATCH CHEMICAL DIVISION

Entrada Industries, Inc. - Salt Lake City, Utah 84106

MORGRO

BENOMYL*
SYSTEMIC FUNGICIDE

ROSES · ORNAMENTALS · FLOWERS · TURF

CAUTION:

**KEEP OUT OF REACH OF CHILDREN
SEE RIGHT PANEL FOR OTHER CAUTIONS**

EPA Reg. No. 168-476

NET CONTENTS: 2 OZ.

CAUTION: MAY IRRITATE EYES, NOSE, THROAT AND SKIN. Avoid breathing dust, contact with skin, eyes and clothing. Wash thoroughly after using. **KEEP AWAY FROM FIRE OR SPARKS.** IN CASE OF CONTACT, flush skin or eyes with plenty of water. IF CONTACTED IN EYES, get medical attention.

IMPORTANT: Do not allow BENOMYL to get wet during storage. Moisture can reduce effectiveness. Keep container closed when not in use. **DO NOT REUSE CONTAINER.** Bury when empty.

NON-WARRANTY: Manufacturer or Seller makes no warranty, express or implied concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mis-handling or any application not specifically described on the label.

ACTIVE INGREDIENT: Benomyl
[Methyl (1-butylcarbamoyl)-2-benzimidazolecarbamate] 50%
INERT INGREDIENTS: 50%

*PATENT PENDING

ACCEPTED
DEC 8 1972
**UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 168-476**
TO ATTACHED COMMENTS.

168-476