

ACCEPTED
 Aug 21, 1973
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE ACT
 FOR ECHO
 168-488

Optional: For use as a pool-side surface disinfectant:

After cleaning surface, use 1 oz. per 2 gallons water for disinfecting. At these dilutions SANITAL will also deodorize surfaces efficiently.

SANITAL will not harm pool equipment. SANITAL is compatible with most commonly used swimming pool chemicals, but should not be mixed with soap or other cleaners or pre-mixed with other water-treating chemicals. SANITAL works best when added by itself directly to pool water.

DANGER: KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbituate drug.

This product is toxic to fish. Do not discharge treated effluent from swimming pools where it will drain into lakes, ponds, streams or public water.

Rinse empty container thoroughly with water and discard.

NON-WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.



SANITAL

DISINFECTANT AND SANITIZER

ACTIVE INGREDIENT:

n-Alkyl (50% C₁₄, 40% C₁₂, 10% C₁₆)

dimethyl benzyl ammonium chloride 10.00%

INERT INGREDIENTS: 90.00%

TOTAL 100.00%

WARNING: KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR OTHER WARNINGS

EPA Reg. No. 168-488

NET CONTENTS: ONE GALLON



SANIT
HARD WATER
 grains) as ca

GENERAL DI
 containing 1
 2 gallons of

DISINFECTIC
 razors for 10

DISINFECTIC
 10 minutes.

COLD DISINI
 1 gallon wat
 rosion inhibi

SANITIZATIO
 thoroughly, t
 mended dilu

FOOD PROC
 tizing solutio

SWIMMING

1. In freshly
algae gro
2. If algae gi
monly fou
add 1 gal
3. If the poo
days to m
4. If pool wa
mine the
tain 2 to



SANITAL

DISINFECTANT AND SANITIZER

ACTIVE INGREDIENT:

Alkyl (50% C₁₄, 40% C₁₂, 10% C₁₆)

dimethyl benzyl ammonium chloride 10.00%

INERT INGREDIENTS: 90.00%

TOTAL 100.00%

WARNING: KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR OTHER WARNINGS

8-488

NET CONTENTS: ONE GALLON



SANITAL DISINFECTANT & SANITIZER SHOULD BE DILUTED BEFORE USE.

HARD WATER TOLERANCE: SANITAL is recommended for use as a sanitizer in waters up to 550 ppm hardness (32 grains) as calcium carbonate when tested by the A.O.A.C. Germicidal and Detergent Sanitizers Official Method.

DIRECTIONS:

GENERAL DISINFECTANT AND SANITIZATION in hospitals, public rooms and at home. After cleaning apply solution containing 1 oz. per 1 gallon of water for hospital disinfectant. For a disinfecting solution in homes, use 1 oz. per 2 gallons of water.

DISINFECTION OF BARBER SHOP AND BEAUTY PARLOR EQUIPMENT: Clean thoroughly then soak combs and razors for 10 minutes in a solution of 1 oz. per 1 gallon of water.

DISINFECTION OF RUBBER GOODS: Clean thoroughly, then soak in solution diluted 1 oz. per 1 gallon of water for 10 minutes.

COLD DISINFECTION OF SURGICAL INSTRUMENTS: Clean thoroughly, rinse and immerse in solution of 1 oz. per 1 gallon water for 10 minutes. 1 tablespoon of sodium nitrite should be added to each gallon of solution as a corrosion inhibitor.

SANITIZATION OF GLASSWARE, SILVERWARE, DISHES, COOKING UTENSILS: Clean with suitable detergent, rinse thoroughly, then dip in a sanitizing rinse made with 1 oz. in 4 gallons water. Air Dry. Do not use a towel. (Recommended dilution should conform with local health department regulations.)

FOOD PROCESSING EQUIPMENT: Use 1 oz. per 4 gallons water. Clean equipment thoroughly, then apply sanitizing solution. Allow equipment to drain and air dry. No potable water rinse is necessary before reuse.

SWIMMING POOL ALGAECIDE

1. In freshly filled pools or pools showing no visible algae growth, add 1 gallon per 25,000 gallons water to prevent algae growth.
2. If algae growth is observed, add 1 gallon for each 10,000 gallons water to control those algae species most commonly found in swimming pools. When existing growth is dead, clean pool, drain and refill if necessary, then add 1 gallon per 25,000 gallons water.
3. If the pool water treated as in (1) or (2) above is recycled, add 1 gallon per 50,000 gallons water every 5 to 7 days to maintain between 2 to 5 parts per million of active ingredient. See serviceman for test kit.
4. If pool water treated as in (1) or (2) above is not recycled or not filtered, it should be tested periodically to determine the active ingredient content. When necessary, 1 gallon per 50,000 gallons water should be added to maintain 2 to 5 parts per million of active ingredient.