



2

in

1

*July 1972*  
*10/2/72*

**weed and feed**



# **20-10-5 PLUS TRIMEC®**

**HERBICIDAL CONCENTRATE**

**KILLS DANDELION, KNOTWEED,  
CHICKWEED, PLANTAIN, HENBIT,  
SPURGE (others on back panel)**

**TREATS  
5,000 SQUARE FEET  
NET WEIGHT: 20 LBS.**

**ACTIVE INGREDIENTS:**

•2,4-dichlorophenoxyacetic acid . . . . .	0.73%
••2-(2-methyl-4-chlorophenoxy) propionic acid . . . . .	0.31%
•••Dicamba (3,6-dichloro-o-anisic acid) . . . . .	0.06%
<b>INERT INGREDIENTS:</b> . . . . .	<b>98.90%</b>
(Relative to herbicidal content) <b>TOTAL</b> . . . . .	<b>100.00%</b>

Equivalent to: •1.266 lbs. per acre  
••0.534 lbs. per acre  
•••0.113 lbs. per acre

TRIMEC® is a registered trademark of FBI-Gordon Corporation, U.S.  
Patent No. 3,284,188

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN  
SEE OTHER CAUTIONS ON BACK PANEL**

**EPA Reg. No. 168-492**



Manufactured by

**WASATCH CHEMICAL DIVISION**

1679 SOUTH SIXTH WEST / P.O. BOX 8128 / SALT LAKE CITY, UTAH 84108

# MORGRO 2 in 1 weed and feed

**20-10-5 PLUS TRIMEC® HERBICIDAL CONCENTRATE**

## GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N) . . . . .	20.00%
6.1% Ammoniacal Nitrogen from Ammonium Sulfate and Diammonium Phosphate	
13.9% Organic Nitrogen from Urea	
AVAILABLE PHOSPHORIC ACID (P <sub>2</sub> O <sub>5</sub> ) . . . . .	10.00%
Derived from Diammonium Phosphate	
SOLUBLE POTASH (K <sub>2</sub> O) . . . . .	5.00%
Derived from Muriate of Potash	
IRON (Fe) Derived from Iron Sulfate . . . . .	1.00%

## DIRECTIONS FOR USE

**WHEN TO APPLY:** For **BEST RESULTS** apply in **spring** or **early fall** when weeds are growing vigorously, after a heavy rain.

Use **2-in-1 WEED & FEED** on Blue Grass, Fescue, Bermuda, and Zoysia lawns.

**2-in-1 WEED & FEED** is a fast, long-lasting fertilizer that produces a thick, green turf while destroying weeds.

**AMOUNT TO USE:** Covers 5,000 sq. ft. Apply evenly with lawn spreader in either one-pass or two-pass application, criss-crossing the area at half-the-rate each time.

FOR MAXIMUM WEED KILLING RESULTS, do not rake, mow or water lawn for at least 24 hours after spreading.

Lawns heavily overrun by weeds and in poor condition may require more than one application. A second application may be made in 30 days.

## SPREADER SETTINGS

MODEL	Jackson	Lawn Beauty	Cyclone Morgro	Scotts	Sears Push Type
SETTING	8	7	3	4	8

Above spreader settings are approximate. CHECK YOUR SPREADER on 250 sq. ft. and adjust to apply at 1 lb. per 250 sq. ft. Use normal rate of walk when applying to avoid error.

### 2-In-1 WEED & FEED controls the following weeds:

Bedstraw	Ground Ivy	Peppergrass	Speedwell
Black Medic	Heal-all	Pigweed	Spurge
Buckhorn	Henbit	Plantains	Thistle
Burdock	Knotweed	Poison Ivy	Wild Carrot
Chicory	Lambsquarter	Poison Oak	Wild Garlic
Chickweed	Lespedeza	Purslane	Wild Lettuce
Clover	Mallow	Ragweed	Wild Onion
Dandelion	Morning Glory	Sheep Sorrel	Yarrow
Dock	Oxalis	Shepherd's Purse	