

DIRECTIONS FOR UDDER WASHING AND SANITIZING

1. For each cow to be milked, place one clean towel in plastic bucket containing fresh, warm 25 ppm titratable iodine solution (½ oz. TAME to a 10-quart pail).
2. Wash udder; discard towel in second bucket.
3. Examine each quarter with strip cup.

DANGER

KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. Rinse empty container thoroughly with water and discard it.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

097504



WASCO

TAME

**DETERGENT AND SANITIZER FOR USE ON DAIRY
FOR UTENSILS • EQUIPMENT • UDDERS**

**DANGER: KEEP OUT OF REACH OF CHILDREN
SEE LEFT PANEL FOR FIRST AID & ADDITIONAL PRECAUTIONS**

ACTIVE INGREDIENTS:

alpha-(p-Nonylphenyl)-omega-hydroxypoly
(oxyethylene)-iodine complex
(Providing 1.75% titratable iodine)
Phosphoric Acid

INERT INGREDIENTS

TOTAL

EPA Reg. No. 168-494

U.S. Patents: 3,028,299; 2,977,315, 2,931,777

NET CONTENTS



Manufactured by

WASATCH CHEMICAL

1009 SOUTH SEVENTH WEST, P.O. BOX 10, SALT LAKE CITY, UTAH 84119

open titratable

Protect eyes
thoroughly

minutes. For eyes,

these are not

gastric lavage.

ACCEPTED
168-494
10 FEB 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT
POISON REGISTERED
UNDER NO.

SANITIZES EQUIPMENT
FAST ACTING • EASY-TO-USE

DIRECTIONS FOR CLEANING UTENSILS AND EQUIPMENT

1. Scrape and prewash dishes, glasses, utensils, and equipment whenever possible.
2. Wash with TAME solution prepared by using 1 oz. to each 5 gallons of warm water.
3. Rinse with potable water.
4. Sanitize in a solution of 1 oz. TAME to 5 gallons of water. Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on rack or drainboard to air dry.

Change washing and sanitizing solutions when the brown iodine color disappears. Make fresh solution each time used.

TABLE OF DILUTION

One-ounce TAME to five U.S. Gallons equals 25 ppm titratable iodine.

NON-WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described and recommended on the label.

EPA Est. 168-UT-1