

USE DIRECTIONS

COLD DISINFECTION

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ ounce SANITAL II per gallon of water for 10 minutes.

SWIMMING POOL ALGAEICIDE

Initial Dose: 1 gallon SANITAL II to 50,000 gallons of water.
Maintenance Dose: 1 quart SANITAL II to 50,000 gallons of water. Initial dose is used when filling pool at beginning of swimming season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat this treatment procedure as necessary. When pool is free of visible algae, use the recommended maintenance dose.

Maintenance dose should be added at 3-5 day intervals. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently.

If make-up water is added at the rate of more than 10% per week of pool capacity, add 1 quart of SANITAL II for each 10,000 gallons of fresh make-up water added per week.

When swimming season is over, add 1 gallon of SANITAL II to 50,000 gallons of water in treating water left in pool. This dose helps provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to use the following season.

When SANITAL II is used as directed, it will kill and prevent development of *Chlorella pyrenoidosa* (green), *Phormidium mundatum* (black algae), *Phormidium retzii* (Square D algae), *Lyngba versicolor* (blue-green) and other swimming pool algae.

SANITAL II is compatible with most commonly used swimming pool chemicals, but should not be mixed with soap or other cleaners, or premixed with other water-treating chemicals. SANITAL II works best when added by itself directly to pool water. SANITAL II at use dilution is non-corrosive to inanimate surfaces.

HOUSEHOLD

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For schools, homes, locker rooms, garbage pails, sink tops, corridors, classrooms, offices and shower stalls. Apply solution with mop or cloth. Sanitize with 1 ounce SANITAL II to 4 gallons of water. Disinfect with 3 ounces SANITAL II to 5 gallons of water. Kills *Staphylococcus aureus* and *Salmonella choleraesuis*. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

HOSPITALS

For floors, walls and other hard surfaces. In nursing homes, institutions, as well as sink tops, garbage pails, telephones and restrooms, use SANITAL II at 3½ ounces per 5 gallons of water.

Apply with cloth, mop, or sponge. At this level SANITAL II will be effective against *Pseudomonas aeruginosa* PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

FOOD PROCESSING EQUIPMENT

For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce SANITAL II per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, NO POTABLE WATER RINSE is required.

DAIRIES

To sanitize dairy equipment such as tanks, lines, pails and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce SANITAL II to 4 gallons of water (200 ppm). At this level, NO POTABLE WATER RINSE is required. Follow recommendations of local Health Board.

EPA Est. 168-UT-1



WASCO

BRAND

097041

DISINFECTANT • DEODORIZER • SANITIZER

SANITAL II

SANITAL II IS A CONCENTRATE AND SHOULD BE DILUTED BEFORE USING.

**DANGER: KEEP OUT OF REACH
OF CHILDREN**

SEE OTHER PRECAUTIONARY AND FIRST AID STATEMENTS
ON RIGHT SIDE PANEL

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides
n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
ethylbenzyl ammonium chlorides

INERT INGREDIENTS:

TOTAL 100%

NET CONTENTS: ONE GALLON

EPA Reg. No. 168-504



Manufactured by

WASATCH CHEMICAL DIVISION

1979 SOUTH SEVENTH WEST / P.O. BOX 6219 / SALT LAKE CITY, UTAH 84141



DISINFECTANT • DEODORIZER • SANITIZER

SANITAL II

SANITAL II IS A CONCENTRATE AND SHOULD BE DILUTED BEFORE USING.

DANGER: KEEP OUT OF REACH OF CHILDREN
 PRECAUTIONARY AND FIRST AID STATEMENTS ON RIGHT SIDE PANEL

ACTIVE INGREDIENTS:	
n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides	5.00%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides	5.00
INERT INGREDIENTS:	90.00
TOTAL	100.00%

NET CONTENTS: ONE GALLON

168-504



USE DIRECTIONS

RESTAURANT AND BAR RINSE

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce SANITAL II to 4 gallons of water. NO TERMINAL POTABLE WATER RINSE REQUIRED.

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against Influenza A₂, Herpes Simplex, Adenovirus type 2, and Vaccinia Viruses.

DANGER

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food. Rinse empty container thoroughly with water and discard.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public waters. Apply this product only as specified on this label.


FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN:

Probable mucosal damage contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

NON-WARRANTY: Manufacturer or Seller makes no warranty, expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Manufacturer or Seller is not liable for any injury or damage caused by this product due to misuse, misbranding or any application not specifically described and recommended on the label.



Comments

ACCEPTED WITH COMMENTS