



CAPTAN 300

UNDYED FLOWABLE SEED PROTECTANT

Agricultural Fungicide - Liquid Suspension

ACTIVE INGREDIENTS:	BY WT.
CAPTAN: N-(1-Trichloromethyl-2-hydroxyethyl)-2,6-dimethyl-4-pyridinecarboxamide	28.4%
Related Derivatives	1.6%
INERT INGREDIENTS	70.0%
	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Statement of Practical Treatment

- *IN CASE OF EYE CONTACT:** Immediately flush eyes with running water for at least 15 minutes. Call a Physician.
- *IN CASE OF SKIN CONTACT:** Wash the affected area with soap and warm water.

See side panel for additional precautionary statements.

STORE IN A COOL, DRY PLACE
STIR THOROUGHLY TO MIX CONTENTS
BEFORE USING.

5 GAL. NET

Manufactured for

AGRICULTURAL PRODUCTS COMPANY, Ft. Worth, TX 76112

SEED TREATMENT DOSAGE RATES FLUID OUNCES per HUNDRED POUNDS of seed

The Following Table gives the amounts of this product to use on various seeds.

SEED	DOSAGE RATE
Alfalfa, Clover, Lespedeza, Trefoil	6.5 - 10.7 fl. oz. 100 lb.
Beans (snap dry, crabeau)	2.5 - 4.0 fl. oz. 100 lb.
Beets (garden)	10.0 - 15.5 fl. oz. 100 lbs.
Cereal Grains (for seed & seedlings only - not for burr smut)	
Barley, rye	2.0 - 4.0 fl. oz. 100 lb.
Oats	2.5 - 5.2 fl. oz. 100 lb.
Wheat	1.5 - 5.2 fl. oz. 100 lb.
Cole Crops (Broccoli, Brussels sprouts, Cabbage, Cauliflow)	1.3 - 2.6 fl. oz. 100 lb.
Corn	
Field Corn	1.5 - 3.0 fl. oz. 100 lb.
Sweet Corn	2.6 - 5.2 fl. oz. 100 lb.
Cotton	
And de-linted cotton	2.5 - 6.5 fl. oz. 100 lb.
Reginned and Fuzzy cotton	3.5 - 9.0 fl. oz. 100 lb.
Cucumbers (Mustard, Radishes, Rape, Turnips)	1.3 - 2.6 fl. oz. 100 lb.
Flax	2.6 - 4.2 fl. oz. 100 lb.
Grasses	6.5 - 13.8 fl. oz. 100 lb.
Milo	5.0 - 8.2 fl. oz. 100 lb.
Muskmelon, Watermelon, Pumpkin, Squash	1.7 - 2.6 fl. oz. 100 lb.
Peanuts (unshelled)	6.0 - 7.8 fl. oz. 100 lb.
Peas	2.5 - 3.5 fl. oz. 100 lb.
Rice	3.0 - 7.8 fl. oz. 100 lb.
Sorghum (thru-hess)	2.2 - 5.8 fl. oz. 100 lb.
Soybeans	2.0 - 3.2 fl. oz. 100 lb.
Sonch	5.0 - 7.8 fl. oz. 100 lb.
Sugar Beets (West East)	5.0 - 9.3 fl. oz. 100 lb.
Surflowers	10.0 - 15.7 fl. oz. 100 lb.
Swiss Chard	2.6 - 5.2 fl. oz. 100 lb.
Tomato, Cucumbers, Cantaloupe	10.0 - 16.0 fl. oz. 100 lb.
	2.5 - 4.0 fl. oz. 100 lb.

ACCEPTED

MAY 27 1981

Use lower rates shown above when method of treating produces consistently even coverage of seed and when favorable planting weather prevails. Use higher rates when weather conditions are unfavorable or uncertain or where method of treating may not consistently produce even coverage of each seed.

STORAGE AND DISPOSAL:

- 1. PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping prohibited. Do not reuse empty container.
- 2. PESTICIDE DISPOSAL:** Pesticide, spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or Local procedures under the Resource Conservation and Recovery Act.
- 3. CONTAINER DISPOSAL:** Dispose of containers according to approved Federal, State or Local procedures under the Resource Conservation and Recovery Act. If container cannot be reused, dispose of in a sanitary land fill or by incineration if permitted by State and Local Authorities.
- 4. GENERAL:** Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

NOTICE-READ CAREFULLY

CONDITIONS OF SALE: Seller warrants that this product is as described and is fit for the uses and purposes intended by the buyer and is not adulterated or misbranded.

WARRANTY LIMITATION: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label, subject to the inherent risks referred to below. Seller makes no other express warranties. **THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY** and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with this use. Buyer assumes all risks associated with use or application of this product contrary to use instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY: In no case shall Seller be liable for special, indirect or consequential damages resulting from the use or handling of this product and not claim for any loss greater in amount than the purchase price of the product in respect of which such damages are claimed.

1977 EPA Reg. No. 13789-21

EPA Est. 43789-TX-01

BO-RID V-4 LIQUID WEED KILLER

ACCEPTED
JUN 4 1981
U.S. Federal Insecticide, Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 179-86

GENERAL INFORMATION

BO-RID "V-4" Liquid Weed Killer is a solution to be applied as a spray on non-cropland and, except for Johnsongrass, is particularly effective for control of most perennial weeds. "V-4" is an effective general herbicide for the control of many annual and biennial weeds. It is absorbed through the foliage and usually exhibits a high degree of contact activity. The degree and duration of effect will vary with the amount of chemical applied, rainfall, temperature, and other conditions.

WATER DISPERSIBLE LIQUID • 1 GAL. CONTAINS .04 LBS. HEXAZINONE

ACTIVE INGREDIENT
Hexazinone [3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H) dione]
INERT INGREDIENTS

EPA Est. 00179-KO1

EPA Reg. 179-89

Keep out of reach of children

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION: CAUSES EYE DAMAGE. FLAMMABLE.

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Environmental Hazards

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

IMPORTANT—Injury to or loss of desirable trees or other plants may result from failure to observe the following. Do not apply except as recommended or direct or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Keep from contact with fertilizers, insecticides, fungicides, and seeds.

Thoroughly clean all traces of V-4 from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzles, tips and other parts. Clean these parts separately.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Net Contents, 10,000 gallons

Manufactured for

THE R. H. BOGLE COMPANY

ALEXANDRIA, VA

JACKSONVILLE, FLA MEMPHIS, TENN

ALVA, OKLA

NOTICE OF WARRANTY

Bogle warrants that this product conforms to the chemical description on the label thereon and is reasonably fit for purposes stated on such label only when used in accordance with directions on label under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Ineffectiveness or other unintended consequences may result because of such factors as soil conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bogle. All such risks shall be assumed by the Buyer. **BOGLE MAKES NO WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

DIRECTIONS FOR USE

It is in violation of Federal law to use this product in a manner inconsistent with its label.

Apply "V-4" as a postemergence spray during active plant growth. Do not apply when weeds are dormant or semi-dormant as the treatment may not be effective. Before spraying, calibrate to determine quantity of "V-4" necessary to uniformly cover measured area to be treated. Use amount of "V-4" to assure coverage.

GENERAL WEED CONTROL—Non-Cropland Areas—Railroad, Highway, Utility and Pipeline Ways, Petroleum Tank Farms, Storage Areas, Industrial Plant Sites, Drainage Ditchbanks.

Use the higher levels of the recommended dosage ranges under the following conditions: species; on fine-textured soils (clay, clay loam); on soils containing more than 5% organic matter; and, on coarse textured soils (sand, loamy sand, gravelly soils) where rainfall exceeds 40 inches per year. Use the lower levels of the dosage range if the growing season is short.

Apply 150 gals. to 300 gals. per acre as a foliar spray for season-long control of many annual and perennial weeds including Bahiagrass*, Bermudagrass*, bluegrass, broomsedge, camphor thistle, chickweed, clovers, dewberry, dog fennel, fescue*, fingergrass, foxtail, guineagrass, honeysuckle, lantana, marehail*, natal grass, plantain, prickly lettuce, ragweed, smartweed, wild rice, wild soybean, wild blackberry and wild carrot.

For contact kill and short term control (up to 3 months) of the above weeds, apply 50 gals. per acre as a foliar spray. Application at 50 gals. to 125 gals. per acre also provides contact kill and control of many annual, biennial and perennial weeds including barnyard grass, bindweed, pigweed, bromegrass, buffalograss*, burdock, cocklebur, crabgrass, crown vetch, curly dock, dogbane*, fiddleneck, filaree, fleabane, goatsbeard vine, goldenrod, kochia, lespedeza, mustard, nutsedge*, orchardgrass*, oxalis, paragrass, pigweed, purslane, quackgrass, red root, spurge, trumpet creeper*, wild oats, wild parsnip and wild star thistle.

Note: Do not use on right-of-ways or other sites where marketable timber or other desirable vegetation is adjacent to the treated area. Do not apply to plants standing in water. Do not use in treated ditches for irrigation.

*Considered hard to kill.

Notice to Buyer: Purchase of this material does not confer any rights under patents of the United States.

BOGLED V-4 WEED KILLER

ACCEPTED
JUN 4 1981
U.S. Federal Insecticide, Fungicide, and Rodenticide Act
as amended for the pesticide
registered under
EPA Reg. No. 179-81

GENERAL INFORMATION

BOGLED V-4 Liquid Weed Killer is a solution to be applied as a spray on non-cropland areas.

"V-4" is an effective general herbicide for the control of many annual and biennial weeds, woody vines, and, except for Johnsongrass, is particularly effective for control of most perennial weeds. "V-4" is readily absorbed through the foliage and usually exhibits a high degree of contact activity. The degree of control and duration of effect will vary with the amount of chemical applied, rainfall, temperature, soil texture and other conditions.

NOTICE OF WARRANTY

Bogle warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bogle. All such risks shall be assumed by the Buyer. **BOGLE MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE**

DIRECTIONS FOR USE

It is in violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply "V-4" as a postemergence spray during active plant growth. Do not apply when vegetation is dormant or semi-dormant as the treatment may not be effective. Before spraying, calibrate equipment to determine quantity of "V-4" necessary to uniformly cover measured area to be treated. Measure proper amount of "V-4" to assure coverage.

GENERAL WEED CONTROL—Non-Cropland Areas—Railroad, Highway, Utility and Pipeline Right-of-Ways, Petroleum Tank Farms, Storage Areas, Industrial Plant Sites, Drainage Ditchbanks.

Use the higher levels of the recommended dosage ranges under the following conditions: on hard-to-kill species; on fine-textured soils (clay, clay loam); on soils containing more than 5% organic matter or carbon; and, on coarse-textured soils (sand, loamy sand, gravelly soils) where rainfall exceeds 40 inches per year. Use the lower levels of the dosage range if the growing season is short.

Apply 150 gals. to 300 gals. per acre as a foliar spray for season-long control of many annual, biennial and perennial weeds including Bahiagrass*, Bermudagrass*, bluegrass, broomsedge, camphorweed, Canada thistle, chickweed, clovers, dewberry, dog fennel, fescue*, fingergrass, foxtail, guineagrass, heath aster, honeysuckle, lantana, marehail*, natal grass, plantain, prickly lettuce, ragweed, smutgrass, Spanish needle, vaseygrass, wild blackberry and wild carrot.

For contact kill and short term control (up to 3 months) of the above weeds, apply 50 gals. to 125 gals. per acre as a foliar spray. Application at 50 gals. to 125 gals. per acre also provides contact kill and short term control of many annual, biennial and perennial weeds including barnyard grass, bindweed*, bouncing bet*, bromegrass, bufflegum*, burdock, cocklebur, crabgrass, crown vetch, curly dock*, dandelion*, dogbane*, fiddleneck, filaree, fleabane, goatshead vine, goldenrod, kochia, lespedeza, milkweed*, mustard, nutsedge*, orchardgrass*, oxalis, paragrass, pigweed, purslane, quackgrass, ryegrass, smartweed, spurge, trumpet creeper*, wild oats, wild parsnip and wild star thistle.

Note: Do not use on right-of-ways or other sites where marketable timber or other desirable plants are immediately adjacent to the treated area. Do not apply to plants standing in water. Do not use water from treated ditches for irrigation.

* Considered hard-to-kill.

Notice to Buyer: Purchase of this material does not confer any rights under patents of countries outside the United States.

1 GAL. CONTAINS .04 LBS. HEXAZINONE

1-methyl-1,3,5-triazine-2,4(1H,3H)-dione] . . . 05%
99.5%

EPA Reg. 179-89.

reach of children

ary Statements

s and Domestic Animals

E DAMAGE, FLAMMABLE.

othing. In case of contact immediately flush
et medical attention if irritation persists

ental Hazards

ot apply where runoff is likely to occur. Do not
equipment or disposal of wastes.

rees or other plants may result from failure to observe
ended or drain or flush equipment on or near desirable
roots may extend, or in locations where the chemical
eir roots. Do not use on lawns, walks, driveways, tennis
to desirable plants. Keep from contact with fertilizers

ication equipment immediately after use. Flush tank
of water after removing nozzle tips and screens (clean

and Disposal

eed by storage or disposal. Pesticide, spray
sed according to label instructions must be
late or local procedures under the Resource

nts: 10,000 gallons.

ufactured for
BOGLE COMPANY
XANDRIA, VA.

LA. MEMPHIS, TENN.
VA, OKLA