

ACCEPTED

May 31, 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. *192-40* SUBJECT
TO ATTACHED COMMENTS.

192-40

Destruxol SPIDER-sMITE is used specifically for the control of various agricultural mites including: European Red Mite, Brown Almond or Clover (Bryobia) Mite, Red Spider Mite, Strawberry Mite, Cyclamen Mite, Pnyet Mite and Citrus Red Mite.

Destruxol SPIDER-sMITE may be used for control of these Mites on most ornamental and flowering plants and shrubs. On sensitive varieties use lower dosage.

Destruxol SPIDER-sMITE when used for the control of these mites on:

ORANGES	CHERRIES	STRAWBERRIES
GRAPEFRUIT	PLUMS	RASPBERRIES
LEMONS	GRAPES	BOYSENBERRIES
APPLES	TOMATOES	CUCUMBERS
PEARS	PEPPERS	MUSKMELONS

should not be applied later than 7 days before harvest. Do not use on Avocados. Do not repeat application on cherries or plums within 30 days.

Destruxol SPIDER-sMITE when used for the control of these Mites on:

WALNUTS	PECANS	FILBERTS
PEACHES	APRICOTS	NECTARINES

should not be applied later than 14 days before harvest.

Destruxol SPIDER-sMITE is compatible with many other insecticides and fungicides but should not be used in combination with highly alkaline materials such as lime or lime-sulphur.

ACTIVE INGREDIENTS

1, 1-bis (chlorophenyl)-2, 2, 2-trichloroethanol	18.5%
INERT INGREDIENTS	81.5%



SPIDER-sMITE

(CONTAINS KELTHANE*)

FOR RED SPIDERS AND OTHER MITES

*Rohm & Haas Trademark

DIRECTIONS FOR USE

Shake well before using. Fill sprayer partly full of water. While agitating contents add Destruxol SPIDER-sMITE at the following recommended dilutions:

- 1 to 2 teaspoonfuls per gallon of water
- 1 to 2 tablespoonfuls per 3 gallons of water
- 2 to 4 ounces per 12 gallons of water
- 1 to 2 pints per 100 gallons of water

Add balance of water. Spray thoroughly, covering upper as well as under side of leaves. Spray when mite insects are observed and repeat as necessary. Destruxol SPIDER-sMITE may be used in most of the popular types of hose-end applicators so long as the manufacturer's proportions are followed. To avoid excessive residue at harvest on food crops do not exceed the recommended dosage, or the time restriction between last application and harvest.

CAUTION

Avoid skin contact, and if contacted wash thoroughly thereafter. Avoid inhalation of mist. Do not take internally. Avoid contamination of food-stuffs. Toxic to fish and wild life. Keep out of lakes, streams and ponds.

INFLAMMABLE. Keep away from open flame. Do not smoke while spraying. Do not spray in the presence of flame.

USDA Reg. No. 192-40

DESTRUXOL CORPORATION, LTD.

TORRANCE, CALIFORNIA