

## DIRECTIONS

Destruxol 10% CHLORDANE DUST  
may be used to control:

**ANTS** — Outside buildings. Dust infested areas thoroughly. In addition, dust ant trails and areas around nests if able to locate. Dust around window and door frames and sills, and openings around piping or ducts where ants might enter premises.

**CAUTION:** This material is not to be used inside homes or households or dairy barns or where milk or milk products are used or handled.

**SOD WEBWORMS OR LAWN MOTH IN LAWNS:** Apply evenly at the rate of about 1 pound to 1000 square feet of lawn area, and sprinkle heavily to wash into top layer of soil. Soil should be moist before application, but do not water excessively for several days following application.

**WIREWORMS, CUTWORMS OR GRUBS IN SOIL** (around ornamental plants or shrubs) Cultivate dust into top four inches of soil at the rate of about 1 1/2 pounds per 100 square feet.

**THRIPS, LEAF MINERS, DIABROTICAS AND CERTAIN CATERPILLARS** on Flowers and ornamental plants. Dust thoroughly when insects appear, in early morning or late afternoon. Repeat treatments as necessary.

**CRICKETS, GRASSHOPPERS** on Flowers and Ornamental Plants: Dust plants and ground area around plants. Repeat treatment as necessary.

**EARWIGS, SOWBUGS (Pillbugs)** around flowers and Ornamental Plants: Cultivate material into top four inches of soil where these insects abound. Apply generously under and around boards, stones and bricks in damp shady areas.

USDA Reg. No. 192-41



# CHLORDANE

## 10% DUST



For Control of ANTS, CUTWORMS, GRUBS,  
SOD WEBWORMS, SOW BUGS, EARWIGS

**SQUASH BUGS, TOMATO HORNWORMS, THRIPS, STINK BUGS, COLORADO POTATO BEETLES, LEAF MINERS AND CERTAIN OTHER GARDEN PESTS:** On garden Vegetables (Beans, Cabbage, Cauliflower, Beets, Carrots, Tomatoes, Potatoes, Broccoli, Brussels Sprouts, Squash, Melons), apply thoroughly when insects appear, preferably in the still air of early morning or late afternoon. Use a hand or power duster and cover insects, as this is a contact insecticide, although a toxic residue will remain for a period of two or three weeks following application. Apply at the rate of about 10 pounds per acre. Repeat applications as necessary. **CAUTION:** Do not use on cabbage or related crops after heads begin to form. Do not apply to cucurbits. Do not apply to beans and related crops after pods begin to form. Limit use on asparagus to post cutting. Do not use on any vegetable crops after edible parts have begun to form. Do not use tops of carrots or other root crops which have been treated with this material for feed or food. Do not apply to feed crops within 30 days of harvest, also, residue remaining at harvest should be removed by washing, brushing or other effective means. Any forage crops treated with this material should not be fed to dairy animals or animals being finished for slaughter.

*Carrots should be washed thoroughly before use. Do not use tops of carrots or other root crops which have been treated with this material for feed or food. Do not apply to feed crops within 30 days of harvest, also, residue remaining at harvest should be removed by washing, brushing or other effective means. Any forage crops treated with this material should not be fed to dairy animals or animals being finished for slaughter.*

*This product is toxic to fish and wildlife if fed out of any body of water. Birds feeding on treated areas may be killed.*

*Active Ingredient: Chloro-4, 7-methanotetrahydroindene, and 4% other related compounds.*

*\*Equivalent to 6% octachloro-4, 7-methanotetrahydroindene, and 4% other related compounds.*

DESTRUXOL CORPORATION, LTD.

TORRANCE, CALIFORNIA