



DESTRUXOL CRAB GRASS KILLER

PINT TREATS 2000 SQ. FT.

DIRECTIONS FOR USE

READ ENTIRE LABEL BEFORE USING

The best time to kill crab grass is in the early summer when the plants are small and before the seed heads start to form. Soil should be moist before treating; therefore water lawn thoroughly and allow to soak in before application. Either tank or hose end sprayer may be used.

For temperatures 80° and over use 3 tablespoonfuls of DESTRUXOL CRAB GRASS KILLER for each gallon of water and spray at the rate of one gallon of dilute spray for each 200 (20 x 10) sq. ft.

On seedlings under 2" tall, one application should be sufficient, but on larger plants a second application may be necessary in 5 to 7 days.

Follow up spot spraying may be necessary for crab grass germinated after treatments.

Treatment would be the same for lawns of Bluegrass, Zoysia, Bermuda and similar grasses.

This product may be used on Dichondra lawns at temperatures under 80° F. Over this temperature some burning may result.

Lawns may be reseeded after crab grass is dead.



CRAB GRASS KILLER

ACTIVE INGREDIENTS:

Octylammonium Methane arsonate 8.00%

Dodecylammonium Methane arsonate 8.00%

INERT INGREDIENTS

84.00%

(All water soluble)

MAY BE USED ON DICHONDRA

CAUTION: Do not spray newly seeded lawn until after the 3rd mowing. Do not use on St. Augustine, Carpet or Centipede grass. Bent and Fescue grasses may be discolored by this product. Water well the day before treatment and do not water again until two days after treatment. Clean sprayer thoroughly after using. Destroy container after using. Keep children and animals off treated areas until product is washed into soil.

WARNING: Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash skin with soap and warm water immediately. Flush eyes with clear water. Avoid contamination of feed and foodstuffs. Do not spray in lakes, streams or ponds, or on desirable vegetation. This product toxic to wildlife. Birds and other wildlife in treated areas may be harmed. Harmful if swallowed.

ANTIDOTE -- If taken internally, induce vomiting and call a physician at once.

KEEP OUT OF REACH OF CHILDREN

USDA Reg. No. 192-59

DESTRUXOL CORPORATION, LTD

TORRANCE, CALIFORNIA