

DESTROYED

PECAN AND OTHER NUTS

CARE: Keep out of reach of children and other animals.

PREPARATION

Apply in any type of sprayer or pump. Thoroughly wet the foliage and fruit. Destroyed insects will be found on the ground. Thoroughly wash the fruit before eating.

Apply when insects first appear and maintain control. Observe all time limitations and do not use for scale control apply at the "crawler" stage.

DO NOT APPLY ON: Birch Trees, Lima Beans, Concord Grapes, Inafon Deluxe Chrysanthemums, Wealthy Apple or varieties ripening at the same time as or before Wealthy, Anderson Yew, Grassula and Canaerti Junipers, Bosc Pears.

Clean and rinse sprayer after spraying. Wash hands and other exposed parts of body thoroughly. If any chemical is spilled on the hands or body wash immediately with soap and water.

APPLES: Apple Aphid, Rust Mites, Coddling Moth, Clover Mite, Tent Caterpillar, San Jose Scale, Bagworms, Mealy Bug. **DO NOT APPLY WITHIN 30 DAYS OF HARVEST.** Do not apply more than 3 applications during fruiting period. Pomace from treated apples should not be fed to livestock.

PEACHES: Plant Bugs, Aphids, Cottony Peach Scale, Japanese Beetles, Oriental Fruit Moth. **DO NOT APPLY WITHIN 30 DAYS OF HARVEST.** Do not apply more than twice during fruiting period. For Lesser Peach Tree Borer thorough wetting of trunks and main branches is important. To control Peach Tree Borer spray all bark areas from ground to scaffold limbs.

PECANS, FILBERTS, WALNUTS: Aphid, Mites, Pecan Nut Case Bearer, Spittlebug. Do not apply more than twice during fruiting period. Do not apply after husk split on Pecans or Walnuts, and on Filberts after nut is visible. Do not graze livestock in treated groves within 21 days after treatment.

GRAPES: Leafhopper, Mealy Bug, Two Spotted Mite, Japanese Beetle. Do not make more than 2 applications during fruiting period and **DO NOT APPLY WITHIN 30 DAYS OF HARVEST.**

PEARS: Stink Bug, Pear Psylla, Plum Curculio, Aphids, Mealy Bug, Forbes Scale, San Jose Scale. Do not make more than 3 applications during fruiting period. **DO NOT APPLY WITHIN 30 DAYS OF HARVEST.**

PLUMS: Mites, Aphids, Bud Moth. Do not make more than 3 applications during fruiting period. DO NOT APPLY WITHIN 21 DAYS OF HARVEST.

PEAS: (Sprout) Bean Beetle, Leaf Roller, Aphids, Leaf Hopper, Bean Beetle. DO NOT APPLY AFTER POD MATURITY. DO NOT APPLY TO WILLOW TWIGS TO LEASTOCK.

PEAS: (Dried) Aphids, Leaf Skeletonizer, Mexican Bean Beetle. DO NOT APPLY WITHIN 10 DAYS OF HARVEST.

PUMPKINS, SQUASH, MELONS: Aphids, Cucumber Beetle, Squash Vine Borer, Pickleworm, Leaf Hoppers. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

PEPPERS AND EGGPLANT: Flea Beetle, Aphids, Mites, Pepper Maggot, Colorado Potato Beetle, Hornworm, White Fly. DO NOT APPLY WITHIN 21 DAYS OF HARVEST. Do not apply more than 3 times during fruiting season.

STRAWBERRIES: Aphids, Meadow Spittlebug, Mites, Lygus Bugs, White Flies, Strawberry Root Weevil, Thrips. Do not apply more than twice in any 35 day period. DO NOT APPLY WITHIN 4 DAYS OF HARVEST.

ORNAMENTAL: Aphids, Thrips, Japanese Beetle, Bagworms, Euonymus Scale, Tent Caterpillars, Pine Leaf Scale, Sowbugs, Lacebugs, Cyclamen Mite, Oak Kermes, Mealy Bugs, Boxwood Leaf Miner, Millipedes, Tarnished Plant Bug.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing or spray mist. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water: for eyes, get medical attention. Wash with soap and water after handling and change clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not use or store near heat or open flames. When container is empty, wash thoroughly and destroy. This product is toxic to fish and wild life. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on label.

Active Ingredients

*Ethion	2.5%
**Encosulfan	0.37
***Malathion	10.73
Petroleum Hydrocarbons	76.13

Inert Ingredients

7.04
100.00

*0,0,0',0'-Tetraethyl S,S'-methylene bisphosphorodithioate
**Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide
***0,0-Dimethyl dithiophosphate o'-diethyl mercaptosuccinate

USDA Reg. No. 192-66