

ACCEPTED
 Sept 7, 1967
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER.
 ED UNDER NO. 192-70 SUBJECT
 TO ATTACHED COMMENTS.

For control of Bean Beetles, Japanese Beetles, Leaf Hoppers, Aphids, Exposed Thrip, Bagworms, Elm Leaf Beetles, Fleas, on vegetables, small fruits, ornaments and pets.

READ ENTIRE LABEL. Use in strict accordance with label directions and cautions.

GENERAL: Apply Destruxol SEVIN 5% Dust whenever insect damage appears. Best results are obtained when applied in early morning or evening. Thoroughly dust upper and lower surfaces of all foliage. **DO NOT** apply in excess, repeat in 7 day intervals as necessary except as noted.

Asparagus — Asparagus Beetle and Apache Cicada. Apply during cutting season, **DO NOT** repeat application within 3 days. **DO NOT** apply within one day of harvest.

Beans — Bean Beetles, Leaf Hoppers, Army Worms, Stink Bugs.

Corn — Corn Earworm, European Corn Borer, Japanese Weevils, Sap Beetles, Flea Beetles, Leaf Hoppers. For insects attacking silk and ears, start when first silks appear and every 2 to 3 days until silks begin to dry. 3 or more applications are often necessary.

Cucumbers — Melons, Pumpkins, Squashworms, Cucumber Beetles, Leaf Hoppers and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if tender foliage is wet or during periods of prolonged high humidity. **DO NOT** use on Watermelons in Florida.

Potatoes — Tomatoes, Eggplant, Peppers, Colorado Potato — Leaf Hoppers, Fall Army Worm, Lace Bug Worms, Stink Bugs.

Head Lettuce — Flea Beetles, Harlequin Bugs, Leaf Hoppers, Army Worms, Stink Bugs. For Cabbage Loopers apply at 5 to 7 day intervals. **DO NOT** apply within three days of harvest.



* SEVIN 5% DUST

**KILLS FLEAS, TICKS, BAGWORMS
AND MANY OTHER INSECTS**

Grapes — Grape Berry Moth, Japanese Beetle, Leaf Rollers, Leaf Folders and Grape Leaf Skeletonizer. Apply just before first brood Leaf Folder larvae emerge from rolls and as needed for Leaf Hoppers.

Strawberries — Spittlebug, Leaf Roller and Strawberry Weevil. **DO NOT** apply within one day of harvest.

Blueberries — Blueberry Maggot, Cherry and Cranberry Fruitworm.

Gaschoppers — Apply Destruxol SEVIN 5% Dust on above crops as needed.

Dogs and Cats — Ticks and Fleas — Apply dust liberally to entire skin area — Avoiding eyes and rub thoroughly into skin. Comb out dead ticks a few hours after treatment. Repeat weekly as needed. **DO NOT** treat kittens under 4 weeks of age.

Dust liberally around all sleeping quarters, kennels and dog houses.

CAUTION

HARMFUL IF SWALLOWED OR INHALED.

Do Not Breathe Dust. Do Not Take Internally.

Avoid Prolonged or Repeated Breathing of Spray.

Skin Contact With Any Insecticide May Be Harmful. Avoid Unnecessary Contact.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N — methylcarbamate) 5%

INERT INGREDIENTS

95%

*U.S. Pat. No. 2903478

USDA Reg. No. 192-70

*Reg. T.M. Union Carbide Corp.

DESTRUXOL CORPORATION, LTD.

TORRANCE, CALIFORNIA