

Pm04 192-70 7/8/98 1/5
(FRONT CONTAINER PANEL)

ACCEPTED

JUL 8 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 192-70

DEXOL

SEVIN 5%

GARDEN DUST

**For Control of Bean Beetles, Leaf Hoppers, Bagworms, Tomato Hornworms,
Armyworm on Vegetables, Small fruits and Ornamentals**

For home garden use only

Active Ingredient:

Carbaryl (1-naphthyl N-methylcarbamate)

5.0%

Inert Ingredients:

95.0%

Total


100.0%

Keep Out of Reach of Children

CAUTION

See back panel for additional precautionary statements

NET (g)



Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

(The following will be used on 1 lb. shaker canisters)

General: Apply whenever insect damage appears. Apply when air is calm to avoid drift and contact with eyes and skin. Best results are obtained when applied in early morning or late evening hours. Wearing household latex or rubber gloves thoroughly dust upper and lower surfaces of foliage with a thin, even film of dust. Avoid excessive applications. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.


(The following will be used with a 41b. Bag.)

General: Apply whenever insect damage appears. Apply when air is calm to avoid drift and contact with eyes and skin. Best results are obtained when applied in early morning or late evening hours. Wearing household latex or rubber gloves apply product with a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Thoroughly dust upper and lower surfaces of foliage with a thin, even film of dust. Avoid excessive applications. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week

VEGETABLES

Asparagus-Asparagus Beetle and Apache Cicada. Apply during cutting season. DO NOT apply within one day of harvest.

Beans-Mexican Bean Beetle, Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvet bean Caterpillar, Flea Beetles, Japanese Beetle, Corn Earworm, Armyworms, Tarnished Plant Bug, Lygus Bugs, Stink Bugs. May be applied up to 3 days before harvest.



Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi- Flea Beetles, Harlequin Bug, Armyworms, Cabbage Caterpillars, Corn Earworm. Do not apply within 3 days of harvest.

Corn-Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetles, Leafhoppers, Flea Beetles. For Corn Borer: Apply thoroughly into the whorls when plant is about knee high. Repeat at weekly intervals until control is achieved. For Corn Earworm: Apply to silks soon as they appear. Re-apply immediately after rain. May be applied up to 2 days before harvest.

Cucumber, Melon, Squash, Pumpkin-Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use on Watermelons in Florida. May be applied up to 3 days before harvest.

Lettuce, Carrots-Flea Beetles, Harlequin Bug, Leafhopper, Armyworms, Cabbage Caterpillars, Corn Earworm, Spittle Bugs, Tarnished Plant Bugs, Lygus Bugs, Six-spotted Leafhopper, Stinkbugs. May be applied up to 7 days before harvest on Carrots. Do not use on Head Lettuce or Leaf Lettuce within 14 days of harvest.


Okra-Corn Earworm, Stink Bugs. May be applied up to 3 days before harvest.

Potato, Peppers, Eggplant-Colorado Potato Beetle, Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs, Stick Bugs, Lace bugs, Tarnished Plant bugs. May be applied up to 7 days before harvest on potatoes and up to 3 days before harvest on peppers and eggplant.

Tomato-Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer. May be applied up to 3 days before harvest.

BERRIES

Strawberries- Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil. May be applied up to 7 days before harvest.



Raspberries, Boysenberries, Blackberries-Japanese Beetles, Leafrollers, Leafhoppers, Orange Tortrix, Raspberry Aphid. **May be applied up to 7 days before harvest.**

Blueberries-Blueberry Maggot, Cranberry Fruitworm, Orange Tortrix. **May be applied up to 7 days before harvest.**

Grapes-Grape Leafhoppers, Grape Leaf Skeletonizer, Red-Banded Leafroller, June Beetles. **May be applied up to 7 days before harvest.**

ORNAMENTALS

To control bagworm, blister beetles, box elderbugs, boxwood leafminer, flea beetles, Japanese beetle, lacebugs, leafhoppers, leafroller, periodical cicada, plant bugs, psyllids, rose slug, rose aphid, scale insects in the crawler stage, and exposed thrip. Apply thoroughly to the infested plants at the first sign of damage and repeat at 7 day intervals or as necessary for good insect control. To obtain scale control, apply in spring and early summer when scale crawlers are present. Avoid excessive applications. Do not use on Boston Ivy or Virginia Creeper.

STORAGE & DISPOSAL


Storage: Store in a cool dry place. Keep in original container and preferably in a locked storage area. **Pesticide Disposal:** Securely wrap original container in several layers of newspaper and discard in trash. **Container disposal:** Do not reuse canister. Discard canister in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

USER SAFETY REQUIREMENTS: Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product. Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. Do not allow children or pets to contact treated area until dust has settled.



FIRST AID

If Swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious person. **If Inhaled:** Remove victim to fresh air. Apply artificial respiration if indicated. **If on Skin:** Remove contaminated clothing. Wash affected areas with soap and water. **If In Eyes:** Flush eyes with plenty of clean water. Get medical attention if irritation persists. **Note to physician:** Carbaryl is a moderate, reversible cholinesterase, inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for use and safety not in strict accordance with directions.

Dexol Industries
9555 James Avenue South
Suite 200
Bloomington, MN 55431

EPA Reg. No. 192-70
EPA Est. No. 192-CA-1
Rev. 11/96

