		192	-9	(C verse before
--	--	-----	----	--------------------

	-	4
niexe le	1	2-28-9

Esse read instructions on reverse before completing form.	Form Appro	·	60. Approval expires 2-28-
SEPA Environmental Protect Washington, DC 20	tion Agency	Registration Amendment Other	OPP Identifier Number
Applicati	ion for Pesticide - Section	n I	
1. Company/Product Number 192-96	2. EPA Product Menege Insecticide-Rodenti	,	Proposed Classification None Restricted
Company/Product (Name) Dexol Malathion Insect Control	PM#		
5. Name and Address of Applicant (Include ZIP Code) Value Garden Supply, LLC. P. O. Box 585 St. Joseph, MO 64502 Check if this is a new address	(b)(i), my product is s to:		th FIFRA Section 3(c)(3) composition and labeling
	Section - II		
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.	()	olication.	NOTIFICATION AUG 2 6 2003
Notification of label change relative to PR Notice 2001-6. This no regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other consistent with the guidance of PR Notice 2001-6 and the requiremay be subject to enforcement action and penalties under section	changes have been made to the labelin to willfully make any false statement to ements of 40 CFR 156.10 and 40 CFR	ng or the confidential state o EPA. I further understar	ement of formula of this nd that if this notification is not
Material This Product Will Be Packaged in:	000001111		
Child-Resistant Packaging Yes No ** Cartification must be submitted Unit Packaging Yes No. per container	Water Soluble Packaging Yes No If "Yes" No. per Package wgt container	2. Type of Contains Metal Plastic Glass Paper Other	
3. Location of Net Contents Information 4. Size(s) Re	etail Conteiner 5.	Location of Label Direct	tions
6. Manner in Which Lebel is Affixed to Product Litho Pape Sten	ograph Other _ or glued icited		
	Section - IV		
1. Contact Point (Complete items directly below for identification)	ion of individual to be contacted, if n	ecessary, to process th	is application.)
Name Carrie Crawford			ne No. (Include Area Code) 4-849ଧ
Cartific f certify that the statements I have made on this form an f acknowledge that any knowlingly false or misleading st both under applicable law.	nd all attachments thereto are true, a	•	8. Detr Application Received (Stamped)
2. Signature	3. Title Regulatory Affairs Manager		
4. Typed Name	5. Date July 21, 2003		

Carrie Crawford 、

Dexol Malathion Insect Control

For Control of Aphids, Mealybugs, Thrips, Scale Insects, Fleas & Other Listed Insects

For Use on Home Ornamentals, Fruits. & Vegetable Gardens

ACTIVE INGREDIENT:

Malathion: 0.0-dimethyl phosphorodithioate of

diethyl mercaptosuccinate50.0% Inert Ingredients:.....<u>5</u>0.0% TOTAL 100.0%

Contains aromatic petroleum derivative solvent.

EPA Reg. No. 192-96 FPA Est. No. 44616-MO-1 Distributed by: Value Garden Supply, LLC.

P.O. Box 585

St. Joseph. MO 64502

(952) 884-6477

www.valuegardens.comNOTIFICATION

KEEP OUT OF REACH OF CHILDREN CAUTION

AUG 2 6 2003

See back/side panel for additional precautionary statements.

NET CONTENTS:

FIRST AID

If swallowed: Immediately call a poison control center or doctor.

Do not induce vomiting unless told to do so by the poison control center or

doctor.

Do not give any liquid to the person.

Do not give anything by mouth to an unconscious person.

If inhaled:

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial

respiration, preferably by mouth-to-mouth, if possible.

·Call a poison control center or doctor for further treatment advice.

If on skin or clothing: Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15-20 minutes.

·Call a poison control center or doctor for treatment advice.

If in eyes:

Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue

rinsing.

·Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For information of this product (including health concerns, medical emergencies or pesticide incidents) call the National Pesticide Information Center at 1-800-858-7378.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin or clothing. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not permit children and pets onto treated areas until this material has dried completely. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life states of amphibians. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

INOTE: THE FOLLOWING DIRECTIONS FOR USE TO BE USED ON PRODUCT PACKAGED AND SOLD WITH HOSE-END SPRAYER ATTACHMENT ONLY.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

[OPTION 1 - HOSE-END SPRAYER ATTACHMENT INSTRUCTIONS]

HOSE-END SPRAYER ASSEMBLY AND USE INSTRUCTIONS: The hose-end sprayer attachment is precalibrated to deliver 2 tsp. per 1 gal. of water. 1) Attach siphonage tube to hose-end sprayer attachment. 2) Screw hose into hose-end sprayer attachment. 3) Remove cap from bottle and insert sprayer attachment. 4) Securely tighten sprayer attachment on bottle. 5) Turn water on. TO START FLOW OF INSECTICIDE, place finger over hole on right side of sprayer attachment. Hold about 3 feet above area to be treated moving at a steady pace to ensure consistent coverage. 6) TO STOP FLOW OF INSECTICIDE, remove finger from hole to break siphon action. NOTE: To ensure adequate coverage. use moderate to high water pressure when applying through hose-end sprayer attachment.

[OPTION 2 - HOSE-END SPRAYER ATTACHMENT WITH NUT INSTRUCTIONS] HOSE-END SPRAYER ASSEMBLY AND USE INSTRUCTIONS: 1) Attach siphonage tube to hoseend sprayer attachment. 2) Remove cap from bottle and insert sprayer attachment. 3) Securely tighten sprayer attachment on bottle. 4) Securely tighten hose into nut on hose-end sprayer attachment. 5) Turn water on. TO START FLOW OF INSECTICIDE, place finger over hole on right side of sprayer attachment. Hold about 3 feet above area to be treated moving at a steady pace to ensure consistent coverage. 6) TO STOP FLOW OF INSECTICIDE, remove finger from hole to break siphon action. NOTE: To ensure adequate coverage, use moderate to high water pressure when applying through hose-end sprayer attachment.

Do not apply to food crops within 3 days of harvest unless otherwise specified.

ORNAMENTALS: (Roses, Camellias, Azaleas, Stocks, Chrysanthemums, Evergreens, and many other flowers and shrubs). Aphids.

NOTE: Injury may occur on tender varieties of plants such as Boston and Maidenhair ferns, pteris and crassula.

FRUITS: (Apples, Pears, Grapes) Aphids. Malathion may cause injury to McIntosh and Cortland varieties of apples, to Bosc Pears and Ribier Grapes. Do not apply to Pears within 1 day of harvest. (Apricots) Codling Moth, Orange Tortix, Terrapin Scale, Soft Brown Scale, Aphids. Do not apply within 7 days of harvest.

(Nectarines) Plum, Curculio, Mites, Parlatoria Scale. Do not apply within 7 days of harvest.

VEGETABLES: (Carrots, Celery, Cucumbers, Endive, Squash, Radishes, Tomatoes) Aphids, Spider Mites. Do not apply within 7 days of harvest on Carrots, Celery, Endive, Radishes; 1 day on Cucumbers. Squash, and Tomatoes. (Lettuce, Spinach) Leafhoppers, Aphids, Mites, Cabbage Looper. Do rict apply on leaf lettuce within 14 days of harvest; head lettuce and spinach within 7 days of harvest; (Onions) Thrips.

FLEAS: In yards, lawns and under houses - Apply Dexol Malathion Insect Control at the rate of 1 pint to cover 3,200 sq. ft. DO NOT apply to pets.

414

[NOTE: THE FOLLOWING DIRECTIONS FOR USE TO BE USED ON PRODUCT PACKAGED AND SOLD WITHOU A HOSE-END SPRAYER ATTACHMENT]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Fill tank sprayer partly full with water. Add recommended amount of Dexol Malathion Insect Control and finish filling with balance of water needed. Spray plants thoroughly when insects appear and repeat applications as necessary. When use in hose-end sprayers, follow manufacturer's instructions. NOTE: Injury may occur on tender varieties of plants such as Boston and Maidenhair ferns, pteris and crassula.

Do not apply to food crops within 3 days of harvest unless otherwise specified.

ORNAMENTALS: (Roses, Camellias, Azaleas, Stocks, Chrysanthemums, Evergreens and many other flowers and shrubs) Aphids - 2 tsp. per 1 gal. water. Red Spider Mites, Mealybugs, Wooly Aphid, Whitefly, Thrips, Tarnished Plant Bug, Four-Lined Leafbug, Bagworm, Rose Leafhopper, Japanese Beetle Adults - 1 tbsp. per gal. water. Scale (Black Scale, Soft Brown Scale, Oyster Shell Scale and certain others) - 1 ½ to 2 tbsp. per gal. water.

FRUITS: (Apples, Pears, Grapes) Aphids - 1 ½ to 2 tsp. per 1 gal. water. Malathion may cause injury to McIntosh and Cortland Varieties of apples, to Bosc Pears and Ribier Grapes. Do not apply to Pears within 1 day of harvest. (Apricots) Codling Moth, Orange Tortix, Terrapin Scale, Soft Brown Scale, Aphids - 1 ½ to 2 tsp. per gal. water. Do not apply within 7 days of harvest. (Peaches) European Red Mite, Two-Spotted Mite, Oriental Fruit Moth, Terrapin Scale, Cottony Peach Scale - 2 tsp. to 1 gal. water. Aphids - 1 ½ to 2 tsp. per 1 gal. water. Do not apply within 7 days of harvest. (Nectarines) Plum Curculio, Mites, Parlatoria Scale - 2 tsp. per 1 gal. water. Do apply within 7 days of harvest.

VEGETABLES: (Broccoli, Brussets Sprouts, Cabbage, Cauliflower, Kale, Beans, Peas, Potatoes)
Aphids - 1 ½ to 2 tsp. to 2 gal. water. Red Spider Mites, Mealybugs, Whiteflies, Thrips, Tarnished Plant
Bugs, Four-Line Leaf Bugs, Bean Leafhoppers, Potato Leafhoppers, Japanese Beetle Adults - 1 tbsp. per
gal. water. Do not apply to Beans within 1 day of harvest. Do not apply to Brussels Sprouts, Cabbage,
Cauliflower or Kale within 7 days of harvest. Use up to harvest on potatoes. (Carrots, Celery,
Cucumbers, Endive, Squash, Radishes, Tomatoes) Aphids, Spider Mites - 1 ½ to 2 tsp. per 1 gal. water.
Do not apply within 7 days of harvest on Carrots, Celery, Endive, Radishes; 1 day on Cucumbers,
Squash and Tomatoes. (Lettuce, Spinach) Leafhoppers, Aphids, Mites, Cabbage Looper - 2 tsp. per 1
gal. water. Do not apply on leaf lettuce within 14 days of harvest; head lettuce and spinach within 7 days
of harvest; (Onion) Thrips - 1 ½ to 2 tsp. per gal. water.

FLEAS: In yards, lawns and under houses - Apply spray containing 5 fl. oz. (10 tbsp.) of Dexol Malathion insect Control per 1 gal. of water at the rate of 1 gallon per 1,000 sq. ft. (1 pint to cover 3,200 sq. ft.) DO NOT apply to pets.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry area away from heat or open flame. Keep in original container and preferably in a locked storage area inaccessible to children and pets.

Disposal: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.