

## DEXOL DIAZINON 12½ E

Dexol Diazinon 12½ E provides positive control of many "hard-to-kill" sucking and chewing insects. It works fast and is residually effective.

### Application Rates for Dexol Diazinon 12½ E:

**Lawns:** Bluegrass, Merion, Fescue, St. Augustine, Dichondra, Bent, and Bermuda) for control of Chinch Bugs, Sod Webworms, Bermuda Mites, Ants, Ticks, Earwigs, and Lawn Billbugs, apply at a rate of 4 tablespoonfuls (2 fl. oz.) in 3 gallons of water per 125 sq. ft., or 16 fl. oz. in 24 gals. of water per 1,000 sq. ft. Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water the lawn thoroughly immediately after treatment.

**Roses and Flowers:** (Roses, Azaleas, Chrysanthemums, Gladioli, Camellias, Carnations, Lilies, and Rhododendrons): for control of Aphids, Mites, Leafhoppers, Thrips, and Dangerous Leaf Miners, apply at a rate of 4 tablespoonfuls (2 fl. oz.) per 3 gallons and wet foliage thoroughly. Spray underside of leaves. Wet plants to the dripping point.

**Trees and Shrubs:** (Oak, Maple, Locust, Holly, Boxwood, Pine, Spruce, and Willow): for control of Aphids, Mites, Leaf Miners, Leafhoppers, Scale Crawlers, and Bagworms, apply at a rate of 4 tablespoonfuls (2 fl. oz.) per 3 gallons of water, and wet leaves and branches to the dripping point.

**Fruits:** Apply at a rate of 4 tablespoonfuls per 3 gallons of water as a full cover spray on the following fruits: (Start spraying after blooming and repeat at 10 to 14 day intervals if needed) -

**Apples and Pears** - for control of Codling Moths; Rg. - Apple, Green Apple, and Woolly Apple Aphids; Mites; San Jose, and Forbes Scale Crawlers. Do not apply within 14 days of harvest.

**Berries:** Raspberries, Dewberries, Loganberries and Raspberries - for control of Leafhoppers; Thrips; Aphids; Two-spotted Mites; and Raspberry Sawflies. Do not apply within 7 days of harvest.

**Citrus Fruits** - for control of Citrus Aphids; Cotton-cushion, and Soft Scale Crawlers; Fruit Tree Leaf Rollers; and Citrus Thrips. Do not apply during bloom. Do not apply within 21 days of harvest.

**Peaches and Nectarines** - for control of Aphids; Brown Mites; Two-spotted Spider Mites, Olive, and San Jose Scale Crawlers; Oriental Fruit Moth; and Peach Twig Borer. Do not apply to peaches within 20 days, or to nectarines within 10 days of harvest.

**Vegetables:** for the following vegetables, 4 tablespoonfuls in 3 gallons of water covers 1,000 sq. ft. of garden: (apply when pests first appear; repeat at 7-10 day intervals or as necessary) -

**Broccoli, Brassica Sprouts, Cabbage, and Cauliflower** - for control of Aphids; Diamondback Moths; and Imported Cabbage-worms. Do not apply within 7 days of harvest.

**Melons and Squash** (Cantaloupes, Casaba, Cucumbers, Honeydew Melons, Marathon, Parson Melons, Watermelons, Summer and Winter Squash) - for control of Aphids; Leafhoppers; Mites; Serpentine Leaf Miners; and Thrips. Do NOT apply to melons and winter squash within 3 days, or to summer squash within 7 days of harvest.

**Tomatoes** - for control of Aphids; Dipterous Leaf Miners; Banded Cucumber Beetles; Fall and Southern Armyworms; and Vinegar Flies. Do not apply within one day of harvest.

**Nuisance Pests in outside areas:** for control of Flies; Ants; Cockroaches; Boxelder Bugs; Clover Mites; Spiders; Ticks; Chiggers; and Earwigs, add 16 fl. oz. per 3 gallons of water and thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling foundations, walls, garbage cans, and into cracks and other places where insects may hide. Do not treat plants or animals with this dosage rate.

(Note: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, and orchids since injury may occur.)

### How to Mix and Apply

**Step #1:** Determine size of spraying job. Measure the area to be sprayed. Make applications when insects first appear, or in advance of an expected infestation. Repeat applications as needed or as noted in the application rates.

**Step #2:** Use clean sprayer. Carefully measure and mix the amount of Dexol Diazinon and water as suggested under application rates.

**Step #3:** Spray as directed. Use medium-fine mist sprays for plants. Thorough coverage is important. Flush sprayer with clean water after each use.

\*Diazinon, a registered trademark of CIBA-GEIGY Corp.

**CAUTION:** Keep out of reach of children. May be harmful if swallowed. May be absorbed through skin. Do not breath spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of food and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals. Do not use, pour, spill or store near heat or open flame. Do not contaminate ornamental fishponds. Do not reuse container. Destroy when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do Not: use in the home.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**Active Ingredients:**

\*O,O-dimethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate . . . . . 12.50%

Aromatic Petroleum Derivative Solvent . . . . . 72.99%

Inert Ingredients: . . . . . 14.51%

Total . . . . . 100.00%

U.S. Patent No. 2,754,243

EPA Reg. No. 192-108-AA

DEXOL INDUSTRIES

TORRANCE, CALIFORNIA 90501  
DALLAS, TEXAS 75214



# DIAZINON 12½ E

## MULTI-PURPOSE SPRAY

- For use on lawns, fruits, vegetables, ornamentals and shade trees.
- Works fast - residually effective
- Contains 12½% Diazinon

DEKAL DIAZINON 12 1/2 E



# DIAZINON 12 1/2 E

## MULTI-PURPOSE SPRAY

- For use on lawns, fruits, vegetables, ornamentals and shade trees.
- Works fast — residually effective
- Contains 12 1/2% Diazinon\*

IRVINE, CALIFORNIA 90501  
DALLAS, TEXAS 75214