

DEXOL VEGETABLE SPRAY

Dexol Vegetable Spray provides positive control of many "hard-to-kill" sucking and chewing insects. It works fast and is residually effective.

Application Rates for Dexol Vegetable Spray. For the following vegetables, 4 tablespoonfuls of Dexol Vegetable Spray in 3 gallons of water covers approximately 1,000 sq. ft. of garden. Apply when pests first appear; repeat at 7-10 day intervals or as necessary.

Beans: For control of Black Bean Aphids, Mexican Bean Beetles, Leaf Miners, and Cucumber Beetles. Do not apply within 7 days of harvest.

Broccoli, Brussel Sprouts, Cabbage, and Cauliflower: For control of Aphids, Diamondback Moth, Imported Cabbageworms. Do not apply within 7 days of harvest.

Carrots: For control of Aphids. Do not apply within 10 days of harvest.

Celery, Kale, Swiss Chard, and Turnip Tops: For control of Aphids, Imported Cabbageworms, Diamondback Moths, Harlequin Cabbage Bug, and Corn Earworms. Do not apply within 12 days of harvest.

Lettuce and Endive: For control of Aphids and Dipterous Leaf Miners. Do not apply within 10 days of harvest.

Melons and Squash (Cantaloupe, Casaba, Crowsfoot, Honeydew Melons, Muskmelons, Persian Melons, Watermelons, Summer and Winter Squash): For control of Aphids, Leafhoppers, Mites, Serpentine Leaf Miners, and Thrips. Do not apply to melons and winter squash within 3 days, or to summer squash within 7 days of harvest.

Potatoes: For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded Cucumber Beetles, Leafhoppers, and Southern Armyworms. Do not apply within 35 days of harvest.

Radishes, Parsnips, and Turnips: For control of Aphids, Flea Beetles, and Dipterous Leaf Miners. Do not apply within 10 days of harvest.

Spinach and Beets: For control of Aphids and Dipterous Leaf Miners. Do not apply within 14 days of harvest.

Sweet Corn: For control of Corn Earworms.

Tomatoes: For control of Aphids, Dipterous Leaf Miners, Banded Cucumber Beetles, Fall and Southern Armyworms, Vinegar Flies. Do not apply within one day of harvest.

How to Mix and Apply

Step #1: Determine size of spraying job. Measure the area to be sprayed. Make applications when insects first appear, or in advance of an expected infestation. Repeat applications as needed or as noted in the application rates.

Step #2: Use clean sprayer. Carefully measure and mix the amount of Dexol Vegetable Spray and water as suggested under application rates.

Step #3: Spray as directed. Use medium-fine mist sprays for plants. Thorough coverage is important. Flush sprayer with clean water after each use.

*Diazinon, a registered trademark of CIBA-GEIGY Corp.

CAUTION: Keep out of reach of children. May be harmful if swallowed. May be absorbed through skin. Do not breath spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of food and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals. Do not use, pour, spill or store near heat or open flame. Do not contaminate ornamental fishponds. Do not reuse container. Destroy when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from you: Cooperative Agricultural Extension Service.

Do Not use in the home.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Active Ingredients:

*O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 12.50%

Aromatic Petroleum Derivative Solvent 72.99%

Inert Ingredients: 14.51%

Total 100.00%

U.S. Patent No. 2,754,243

EPA Reg. No. 192-109-AA

DEXOL INDUSTRIES

TORRANCE, CALIFORNIA 90501

DALLAS, TEXAS 75214



- Provides positive control of many sucking and chewing insects.
- Fast working, residually effective
- Contains Diazinon*



RECORDED

beans. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 35 days of harvest.

Broccoli, Brussels Sprouts, Cabbage and Cauliflower. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 14 days of harvest.

Carrots. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 10 days of harvest.

Collards, Kale, Swiss Chard, and Turnip Tops. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 10 days of harvest.

Lettuce and Endive. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 14 days of harvest.

Melons and Squash (Cantaloupes, Casabas, Greenshaws, Honeydew Melons, Muskmelons, Persian Melons, Watermelons, Summer and Winter Squash). For control of Aphids, Leafhoppers, Mites, Serpentine Leaf Miners, and Tobacco Rootworms. Do not apply within 7 days of harvest.

Potatoes. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 35 days of harvest.

Radishes, Parsnips, and Turnips. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 10 days of harvest.

Spinach and Beets. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 14 days of harvest.

Sweet Corn. For control of Flea Beetles.

Tomatoes. For control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leaf Miners, Banded and Green Beetles, leafhoppers, and Southern Armyworms. Do not apply within 35 days of harvest.

How to Mix and Apply:

1. Mix 1/2 ounce of Vegetable Spray per gallon of water. For control of sucking insects, use 1/4 ounce per gallon of water. For control of chewing insects, use 1/2 ounce per gallon of water.

2. Apply to the undersides of leaves and to the soil around the base of the plant.

3. Repeat applications every 7 to 10 days.

4. Do not apply to plants that are stressed or in bloom.

5. Do not apply to plants that are near water.

6. Do not apply to plants that are near children or pets.

7. Do not apply to plants that are near food.

8. Do not apply to plants that are near water.

9. Do not apply to plants that are near children or pets.

10. Do not apply to plants that are near food.

© 1985 DEXAL CORPORATION
 411 CAS TEL BLVD, #575, 14



VEGETABLE Spray

- Provides positive control of many sucking and chewing insects.
- Fast working, residually effective
- Contains Diazinon*

