

[192-114]

Directions for use and other instructions are printed on the side panel of the container.

DIRECTIONS FOR USE

**SHAKE WELL BEFORE USING.
ALWAYS HOLD UPRIGHT WHILE SPRAYING**

For use on all vegetable gardens. Apply to plants as soon as they appear. Repeat applications as needed. Do not use on plants that are stressed or in bloom.

SPRAY CHART

FOOD CROP	INSECTS CONTROLLED
Beans	Bean Beetles, Hornworms, Cucumber Beetles, Flea Beetles, Aphids, and Mexican Weevils
Cucumbers	Aphids, Cucumber Beetles
Peas	Aphids, Flea Beetles
Peppers	Pepper Weevils, Beetles, and Spiders
Tomatoes	Tomato Pinworms, Aphids, and Mexican Beetles
Watermelons	Watermelon Beetles and Spiders

CAUTION

Do not use on plants that are stressed or in bloom. Do not use on plants that are in contact with other plants.

WARNING

Keep out of reach of children. Do not use in areas where children play. Do not use near water.

DEXOL INDUSTRIES

MAKING USE UP TO ONE BEFORE HARVEST

Dexol



CAUTION: Contents under pressure
CAUTION: Keep out of reach of children
See side panel for other cautions
NET WEIGHT 14 OZS.

Under the Federal Insecticide, and Fungicide Act, registered under EPA Reg. No. 192-114
ACCEPTED
1975

DEXOL Vegetable Garden Insect Spray may be used on the food crops listed on this label up to one day before harvest. Used as directed, DEXOL Vegetable Garden Insect Spray will give quick knockdown and kill of those crawling and sucking insects listed.

DIRECTIONS FOR USE

**SHAKE WELL BEFORE USING.
ALWAYS HOLD UPRIGHT WHILE SPRAYING**

Read Entire Label and Apply Only As Directed on This Label.

For best results, spray in evening or on a cloudy day when air is still. Use slow sweeping motion, holding the dispenser approximately 2 feet from foliage. Spray rows at the rate of one foot per second, ensuring that both sides of the leaf are treated. For heavy growth, spray rows on both sides. Spraying should begin when insects first appear and should be repeated as needed, or after rain. Contact insects directly if possible. To prevent possible plant injury, avoid visibly wetting foliage.

SPRAY CHART

FOOD CROP	INSECTS CONTROLLED
Tomato	Flea Beetle, Hornworm, Colorado Potato Beetle, Stink Bug, Aphid, Spotted Mites, White Flies
Cabbage	Aphids, Cabbage Worm
Squash	Aphids, Stinkbug
Peppers, Egg Plant, Potatoes	Colorado Potato Beetle, Leaf Hoppers, Blister Beetle, Aphids, Flea Beetle
Melons	Cucumber Beetles, Aphids Watermelon Beetle
Sweet Corn	Japanese Beetle, Flea Beetle

CAUTION

Do not apply this spray 24 hours before harvest. Wash vegetables and fruits before eating. Avoid using in hot, bright sunlight.

Harmful if swallowed, inhaled or absorbed through skin. Do not breathe spray and avoid contact with eyes. Wash thoroughly after use.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

ACTIVE INGREDIENTS

Pyrethrins	0.56%
Rotenone	.125%
Other cubic resins	2.50%
Petroleum distillate	2.25%

INERT INGREDIENTS

99.344%

Total - 100.00%

E.P.A. Reg. No. 192-114 AA

E.P.A. Est. 33690 CA-1

DEXOL INDUSTRIES

Torrance, Ca. 90501

Dallas, TX 95214

**VEGETABLE
GARDEN
INSECT SPRAY**

**Controls: APHIDS, TOMATO
HORNWORMS, WHITE FLIES,
CABBAGE WORM, SPOTTED
MITES and LEAF HOPPERS.**

**For use on TOMATOES, CAB-
BAGE, SQUASH, PEPPERS,
EGG PLANT, POTATOES,
MELONS and SWEET CORN**

EXPIRES

Federal Register
1/17/74