

... TOMATOES and VEGETABLES... insects and...

USE ON

Tomatoes - Tomato Hornworm, Tomato Fruitworm, Flea Beetle, Early and Late Blight, ...

Peppers and Eggplant - Flea beetle, Tarnished Plant Bug, Colorado Potato Beetle, ...

APPLICATION MAY BE MADE ON ABOVE VEGETABLES UP TO DAY OF HARVEST

Cucumbers, Squash and Melons - Striped and Spotted Cucumber beetle, Squash Bug, Flea Beetle, ...

Beans - Mexican Bean Flea Beetle, Flea and Japanese Beetles, Armyworms, Stink Bugs, ...

Brussels Sprouts, Cabbage, Cauliflower, Broccoli and Kohlrabi - Imported Cabbage Worm, ...

Carrots - Leafhopper

DO NOT APPLY TO BEANS, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, BROCCOLI AND CARROTS WITHIN 7 DAYS OF HARVEST. DO NOT APPLY TO KOHLRABI AFTER EDIBLE PARTS ARE HALF GROWN.

Sweet Corn - European Corn Borer, dust every 5-7 days from the time eggs are about to hatch (when corn is knee high on most varieties). Corn Earworm, dust silks as soon as silks appear. Make at least 3 applications at 2-3 day intervals. Application of this product to the leaf region of corn during the pollen shed period will seriously reduce the bee population.

MAY BE APPLIED TO CORN UP TO DAY OF HARVEST. DO NOT FEED LUNGE TO DAIRY ANIMALS OR ANIMALS BEING FINISHED FOR SLAUGHTER.

DIRECTIONS

Gently shake dust. Hold 1/4 to 1/2 cup from ground level. Hold plant to be dusted, spray or cluster. Apply a continuous stream of dust above ground level, rate of 1 lb. per 100 sq. ft. of ground. For squash and melons use at rate of 1 lb. per 200 sq. ft. of ground. Repeat application at 7-10 day intervals if necessary.

CAUTION: KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed or inhaled or absorbed through skin, eyes and clothing. Do not breathe dust. Wash thoroughly. CALL FOR PHYSICIAN. First aid kit and antidote is antidotal.

This product is toxic to fish and wildlife. Birds, bees, and other beneficial insects should be kept out of any food, or water. Do not apply where bees are present. Do not apply where small birds, fish, or other animals are present.

This product is highly toxic to bees exposed to it. Protective information may be obtained from the manufacturer.

Do not mix or combine with any cleaning or degreasing agent. Do not use in any place where it may be empty.

DO NOT MIX WITH OTHER PRODUCTS. Do not mix with any other product. Do not mix with any other product. Do not mix with any other product.

ACTIVE INGREDIENTS

... ZNEB ... SEVIN ...

INERT INGREDIENTS

... TOTAL ...

EPA REG. NO. 192-61-1

OF AGRICULTURAL INDUSTRIES

Contains ZNEB and SEVIN

Dexco

TOMATO & VEGETABLE DUST

MAY BE USED ON TOMATOES UP TO DAY OF HARVEST.



KILLS HORNWORMS, ARMYWORMS, MEXICAN BEAN BEETLES, FLEA BEETLES, LEAF HOPPERS and controls DOWNY MILDEW, ANTHRACNOSE, RUST BLIGHT, and other listed plant diseases on tomatoes and vegetables.

KEEP OUT OF REACH OF CHILDREN CAUTION SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS NET WT. 10 OZ.

ACCIDENT FEB 26 1979 Under the EPA as directed by registered under EPA Reg. No. 192-61-1