
211112 REGH192 - 11

### (FRONT PANEL - CONTAINER LABEL)

. . . : :

i

DEXOL



1

DEXA-KLOR INSECT SPRAY

Kills Ants, Fleas, Brown Dog Ticks, Grasshoppers, Imported Fire Ants, Chiggers, Grubs, Spiders, Crickets, & More.

[The following statement will be used for products sold with "PESTS OF LAWNS" directions]:

Covers Up To 2,666 Sq.Ft. of Lawn (1 pint size containers only) Covers Up To 5,332 Sq.Ft. of Lawn (1 quart size containers only; Covers Up To 10,664 Sq.Ft. of Lawn (1/2 gallon size containers only)

[The following statement will be used for products sold with "OUTDOOR CONTROL OF FLEAS & TICKS: IN YARDS" directions only]:

Covers Up To 2,560 Sq.Ft. of Yard (1 pint size containers only) Covers Up To 5,120 Sq.Ft. of Yard (1 quart size containers only) Covers Up To 10,240 Sq.Ft. of Yard (1/2 gallon size containers only)

## (BACK PANEL - CONTAINER LABEL)

### SER ATTACHED PAMPHLET FOR COMPLETE DIRECTIONS FOR USE.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not permit children or pets to go on to treated areas until the spray has completely dried. Do not contaminate ornamental fish ponds. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

## STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING UNLESS UNDER MEDICAL SUPERVISION. Product petroleum distillates which may cause aspiration pneumonia. contains If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated. If On Skin: Remove contaminated clothing. Wash affected areas with soap and water. If In Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists. NOTE Chlorpyrifos is a cholinesterase inhibitor. TO PHYSICIAN: Treat symptomatically. Atropine by injection is an antidote.

## ENVIRONMENTAL HAZARDS


This product is toxic to fish, birds and other wildlife. Do not apply to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of waste. This pesticide is highly toxic to bees exposed to treatment or residues remaining on the direct treated area. information may be obtained Protective from your cooperative Agricultural Extension Service.

#### PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

NOTICE: Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.

DEXOL INDUSTRIES Torrance, CA 90501 EPA Reg. No. 192-142 EPA Est. 192-CA-1



· 7


# (SIDE PANEL - CONTAINER LABEL)

. . :

ł

**DIRECTIONS FOR USE:** Dexol Dexa-Klor Insect Spray can be used in yards, kennels & Dog Houses for effective control of Fleas and Brown Dog ticks. Use also in lawns to control Ants, Mole Crickets, Chiggers, and other listed pests. Read attached pamphlet before applying this product.

.



.

DEXOL

### DEXA-KLOR INSECT. SPRAY

## Kills Ants, Fleas, Brown Dog Ticks, Grasshoppers, Imported Fire Ants, Chiggers, Grubs, Spiders, Crickets, & More.

[The following statement will be used for products sold with "PESTS OF LAWNS" directions]:

Covers Up To 2,666 Sq.Ft. of Lawn (1 pint size containers only) Covers Up To 5,332 Sq.Ft. of Lawn (1 quart size containers only) Covers Up To 10,664 Sq.Ft. of Lawn (1/2 gallon size containers only)

[The following statement will be used for products sold with "OUTDOOR CONTROL OF FLEAS & TICKS: IN YARDS" directions only]:

Covers Up To 2,560 Sq.Ft. of Yard (1 pint size containers only) Covers Up To 5,120 Sq.Ft. of Yard (1 quart size containers only) Covers Up To 10,240 Sq.Ft. of Yard (1/2 gallon size containers only)

Active Ingredient: *Chlorpyrifos (0,0-diethyl 0-{3,5,6- trichloro-2-pyridyl] phosphorothioate) Inert Ingredients:	
Total:	
This product contains petroleum distillates.	
*Dursban - Registered Trademark of Dow Chemic	cal CompanyU.S.,
Patent No. 3,244,586.	• • • • • • • • • • • • • • • • • • • •
Keep Out of Reach of Children	
CAUTION	• •
See back panel for additional precautionary	y statements :::
Net Contents	
	••••
	• • •
ł do statu i	• • •

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Dexol Dexa-Klor Insect Spray provides effective control of listed insects that attack lawns, yards, kennels, and listed ornamentals.

DO NOT APPLY TO VEGETABLE GARDENS OR OTHER PLANTS INTENDED FOR FOOD OR FEED PURPOSES.

[THE FOLLOWING "DIRECTIONS FOR USE" WILL BE USED ON PRODUCT SOLD WITH A HOSE-END SPRAYER ATTACHMENT]:

[OPTION 1 - HOSE-END SPRAYER ATTACHMENT INSTRUCTIONS]:

. :

HOSE-END SPRAYER ASSEMBLY AND USE INSTRUCTIONS: 1) Attach siphonage hose-end sprayer attachment. 2) Screw hose tube to into hose-end sprayer attachment. 3) Remove cap from bottle and insert sprayer attachment. 4) Securely tighten spray attachment on bottle. 5) Turn water on. TO START FLOW OF INSECTICIDE, place finger over hole on right side of sprayer attachment. Hold about 3 feet above area t.o be treated moving at a steady pace to ensure consistent coverage. 6) TO STOP FLOW OF INSECTICIDE, remove finger from hole to break siphon NOTE: To ensure adequate coverage, use moderate to high action. water pressure when applying through hose-end sprayer attachment.

**[OPTION 2 - HOSE-END SPRAYER ATTACHMENT WITH NUT INSTRUCTIONS]** HOSE-END SPRAYER ASSEMBLY AND USE INSTRUCTIONS: i) Attach siphonage tube to hose-end sprayer attachment, 2) Remove cap from bottle and insert sprayer attachment. 3) Securely tighten sprayer attachment on bottle. 4) Securely tighten hose into nut on hose-end sprayer attachment. 5) Turn water on. TO START FLOW OF INSECTICIDE, place finger over hole on right side of sprayer attachment. Hold about 3 feet above area to be treated moving at a steady pace to ensure consistent coverage. 6) TO STOP FLOW OF INSECTICIDE, remove finger from hole to break siphon action. NOTE: To ensure adequate coverage, moderate to high water pressure when applying through hose-end use sprayer attachment.

OUTDOOR CONTROL OF FLEAS & TICKS: In Yards - Apply 1 pint to cover 2,560 sq.ft. (1 quart to cover 5,332 sq.ft. or 1/2 gallon to cover 10,240 sq.ft.) of area. Retreat as necessary to maintain control. Thoroughly water immediately after treatment to wash insecticide into For best results, lawn should be moist treated area. attime of treatment. ..... 1N KENNELS AND DOG HOUSES: DO NOT USE HOSE-END SPRAYER ATTACHMENT PROVIDED WITH THIS PRODUCT: Using an adjustable hose-end sprayer or tank type sprayer, apply at the rate of 10-1/2 fl.oz. peril gal. water to building, resting areas, walls and floors. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. DO NOT treat pets with this product. Remove pets before DO NOT return pets to treated areas until spray has dried spraying. completely. .

LAWNS: For control of Ants, Armyworms, Brown Dog Ticks, PESTS OF Chinch Bugs, Clover Mites, Mole Crickets, Cutworms, Flea Beetles, Earwigs, Fleas, Grasshoppers, Sod Webworms -Chiggers, Dichondra Apply 1 pint to cover 2,666 sq.ft. (1 quart to cover 5,332 sq.ft. or 1/2 gallon to cover 10,664 sq.ft.) of area. Spray when pests first Repeat as necessary. Thoroughly water immediately after appear. treatment except as noted. For best results the lawn should be moist at the time of treatment. One application will usually control Chinch Bugs for up to 8 weeks. For Sod Webworms, delay watering 'or mowing the treated area for 12-24 hours after treatment. For Turfgrass (Hyperodis) Weevil, apply 1 pint to cover 1,333 sq.ft. (1 quart to cover 2,666 sq.ft. or 1/2 gallon to cover 5,332 sq.ft.) of area. Spray in mid-April and again in mid-May and follow recommended use For White Grubs (European Chafer and Japanese directions. Beetle larvae only), apply 1 pint to cover 666 to 1,333 sq.ft. (1 quart to cover 1,333 to 2,666 sq.ft. or 1/2 gallon to cover 2,666 to 5,332 sq.ft.) of area. Spray when grubs are young and actively feeding near the soil surface, usually late July and August. Immediately after spraying, water the treated area with 1/2-2" of water. Consult your local Agricultural Extension Service Specialist for the proper time to treat in your area for Hyperodis Weevil and White Grubs. Damaged and ornamental turf should be fertilized to stimulate regrowth lawns and recovery from insect injury.

PESTS OUT OF DOORS: To control Ants, Brown Dog Ticks, Chiggers, Chinch Bugs, Cockroaches, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Flies, Grasshoppers, Sod Webworms, Millipedes, Mosquitoes, Sowbugs, and Spiders - apply to outside surfaces of buildings, porches, screens, window frames, eaves, patios and garages. Repeat as To help prevent infestations of buildings needed. by Ants, Cockroaches, Crickets, Earwigs, Millipedes and Sowbugs: Apply 1 pint to cover 8,000 sq.ft. (1 quart to cover 16,000 sq.ft. or 1/2 gallon to cover 32,000 sq.ft.) of area. Treat a band of soil 6 - 10 ft. wide around and adjacent to the building and 2 - 3 ft, high on the foundation.

Dexol Dexa-Klor Insect Spray may also be used on the sites listed below. DO NOT use the hose-end sprayer attachment provided with this package. Use an adjustable hose-end sprayer or tank type sprayer unless otherwise instructed below.

PESTS INDOORS: For spot treatment of Ants, Cockroaches, Clover Mites, Crickets, firebrats, Silverfish and Spiders - apply as a coarse spray or with a paint brush to corners, cracks in walls, along baseboards, beneath sinks, stoves and around plumbing. FOR ANTS: Apply to ant trails, around doors and windows. Use a 0.25% spray solution to control light infestations or a 0.50% spray solution to quickly reduce heavy infestations. To make a 0.25% spray solution, Mix 5-1/3 fl.oz. in 1 gal. of water or 1 pt. in 3 gals. water. To make a 0.50% spray solution, mix 11 fl.oz. in 1 gal. of water or 1 qt. in 3 gals. water. Always spot test areas for color fastness before spraying.

. :

IMPORTED FIRE ANTS: Dilute 2 fl.oz. per gal. of water. Apply 1 gal. of solution as a gentle rain to each mound using a sprinkling can. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12.5 sq.ft.). For best results, apply in cool weather, 65-80 F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity and treat all colonies which may not as yet have constructed a mound.

ORNAMENTALS (Flowers, Shrubs, Vines, Shade and Flowering PESTS OF and Evergreens) - NOTE: DO NOT USE ON AZALEAS. CAMELLIAS, Trees POINSETTIAS, ROSE BUSHES OR VARIEGATED IVY BECAUSE OF POSSIBLE INJURY. To control Aphids, Bagworms, Eastern Tent Caterpillars, Fall Webworm, Grasshoppers, Orangestriped Oakworm, Yellownecked Caterpillar, Spittlebugs and Whiteflies: Mix 6 fl.oz. in 9 gals. water. Mites. Spray lower leaf surfaces and infested limbs and trunk. Avoid upper and spraying to the point of excessive run-off. Treat when pests appear repeat at 7-10 day intervals if needed. To control Cutworms, and Leafhoppers, Mealybugs, Mimosa Webworm, Redhumped Caterpillar, Thrips (exposed): Use 12 fl.oz. in 9 gals. water. For control of Ash and Lilac Borers, Oak Pine Needle, Tea and White Peach Scale Crawlers: Use 12 fl.oz. in 12 gals. water. NOTE: 1 fl.oz. = 2 tbsp. = 6 tsp.

STORAGE & DISPOSAL

**STORAGE:** Store in a cool, dry place away from heat or open flame. Keep in original container and preferably in a locked storage area. **DISPOSAL:** Do not reuse empty container or hose-end sprayer attachment. Wrap container and hose-end sprayer attachment in newspaper and discard in trash.

[THE FOLLOWING "DIRECTIONS FOR USE" WILL BE USED ON PRODUCT SOLD <u>WITHOUT A HOSE-END SPRAYER ATTACHMENT</u>]

PESTS OF LAWNS: For control of Ants, Armyworms, Brown Dog Ticks, Chiggers. Chinch Bugs, Clover Mites, Mole Crickets, Cutworms, Dichondra Flea Beetles, Earwigs, Fleas, Grasshoppers, Sod Webworms: Apply 3 fl.oz. per 15 gals. water to cover 500 sq.ft. (1 pint to cover 2,666 sq.ft.) of area. Spray when pests first appear. Repeat as Thoroughly water immediately after treatment necessary. except as best results, the lawn should be moist at noted. For time of One application will usually control Chinch Bugs for up to treatment. 8 weeks. For Sod Webworms, delay watering or mowing the treated area for 12-24 hours after treatment. For Turfgrass (Hyperodis) Weevil, apply 6 fl.oz. per 15 gals. water to cover 500 sq.ft. (1 pint to cover 1.333 sq.ft.) of area. Spray in mid-April and again in mid May and follow recommended use directions. For White Grubs (European Chafer and Japanese Beetle larvae only), apply G-12 fl.oz. per 30 gals. water pint to cover 666 to 1,333 sq.ft.) of area. Spray when grubs (] are young and actively feeding near the soil surface, usually late July and August. Immediately after spraying, water the treated area with 1/2-2" of water. Consult your local Agricultural Extension Service Specialist for the proper time to treat in your area for Hyperodis Weevil and White Grubs. Damaged lawns and ornamental turf should be

fertilized to stimulate regrowth and recovery from insect injury.

PESTS INDOORS: For spot treatment of Ants, Cockroaches, Clover Mites, Crickets, Firebrats, Silverfish and Spiders: Apply as a coarse spray or with a paint brush to corners, cracks in walls, along baseboards, beneath sinks, stoves and around plumbing. FOR ANTS: Apply to ant trails, around doors and windows. Use a 0.25% spray solution to control light infestations or a 0.50% spray solution to quickly reduce heavy infestations. To make a 0.25% spray, mix 5-1/3 fl.oz: in l gal. of water or 1 pt. in 3 gals. water. To make a 0.50% spray, mix 11 fl.oz. in l gal. of water or 1 qt. in 3 gals. water. Always spot test areas for color fastness before spraying.

OF DOORS: To control Ants, Brown Dog Ticks, PESTS OUT Chiggers, Chinch Bugs, Cockroaches, Clover Mites, Crickets, Cutworms, Earwigs, Grasshoppers, Sod Webworms, Millipedes, Mosquitoes, Fleas, Flies, Sowbugs and Spiders - apply 3 fl.oz. in 15 gals. water as a spray  $\circ r$ with a paint brush to outside surfaces of buildings, porches, screens, window frames, eaves, patios and garages. Repeat as needed. To help prevent infestations of buildings by Ants, Cockroaches, Crickets, Earwigs, Millipedes and Sowbugs: Apply 2 fl.oz. in 12 gals. water to cover 1,000 sq.ft. (1 pint to cover 8,000 sq.ft.) of area. Treat a band of soil 6-10 ft. wide around and adjacent to the building and 2-3 ft. high on the foundation.

OUTDOOR CONTROL OF FLEAS & TICKS: In Yards - Apply 1-1/4 fl.oz. per 6 water to cover 200 sq.ft. (1 pint t) cover 2,560 sq.ft.) of gals. area. Retreat as necessary to maintain control. Thoroughly water immediately after treatment to wash insecticide into treated area. In Kennels and Dog Houses - Apply 10-1/2 fl.oz. per l gal. water tobuilding, resting areas, walls and floors. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. DO NOT treat pets with this product. Remove pets before spraying. Do not return pets to treated areas until spray has dried completely.

IMPORTED FIRE ANTS: Dilute 2 fl.oz. per gal. of water. Apply 1 gal. of solution as a gentle rain to each mound using a sprinkling can. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12.5 sq.ft.). For best results, apply in cool weather, 65-80 F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity and treat all colonies which may not as yet have constructed a mound.

PESTS OF ORNAMENTALS (Flowers, Shrubs, Vines, Shade and Flowering Trees and Evergreens) - NOTE: DO NOT USE ON AZALEAS, CAMELLIAS, POINSETTIAS, ROSE BUSHES OR VARIEGATED IVY BECAUSE OF POSSIBLE INJURY. To control Aphids, Bagworms, Eastern Tent Caterpillars, Fall Webworm, Grasshoppers, Orangestriped Oakworm, Yellownecked Caterpillar, Mites, Spittlebugs and Whiteflies: Mix 6 fl.oz. in 9 gals. water. Spray upper and lower leaf surfaces and infested limbs and trunk. Avoid spraying to the point of excessive run-off. Treat when pests appear

BEST AVAILABLE COPY

and repeat at 7-10 day intervals if needed. To control Cutworms, Leafhoppers, Mealybugs, Mimosa Webworm, Redhumped Caterpillar, Thrips (exposed): Use 12 fl.oz. in 9 gals. water. For control of Ash and Lilac Borers; Oak Pine Needle, Tea and White Peach Scale Crawlers: Use 12 fl.oz. in 12 gals. water. NOTE: 1 fl.oz. = 2 tbsp. = 6 tsp.

### STORAGE & DISPOSAL

STORAGE: Store in a cool, dry place away from heat or open flame. Keep in original container and preferably in a locked storage area. DISPOSAL: Do not reuse empty container. Wrap container in newspaper and discard in trash.

## [NOTE: THE FOLLOWING PRECAUTIONARY STATEMENTS WILL BE USED ON PRODUCT' SOLD WITH AND WITHOUT A HOSE-END SPRAYER ATTACHMENT]:

### PRECAUTIONARY STATEMENT'S HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed, inhaled or absorbed through the CAUTION: skin. Avoid breathing vapors or spray mist. Avoid contact with skin, eves Wash thoroughly after handling and before eating or and clothing. Avoid contamination of feed and foodstuffs. Do not permit smoking. the children or pets to go on to treated areas until spray has completely dried. Do not contaminate ornamental fish ponds. Food utensils such as teaspoons or measuring cups should not be used for food purposes after use with pesticides.

### STATEMENT OF PRACTICAL TREATMENT

Call a physician or Poison Control Center immediately. If Swallowed: DO NOT INDUCE VOMITING UNLESS UNDER MEDICAL SUPERVISION. Product contains petroleum distillates which may cause aspiration pneumonia. If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated. If On Skin: Remove contaminated clothing. Wash with soap and water. If In Eyes: affected areas Flush eyes with plenty of water. Get medical attention if irritation persists. NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection is an antidote.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Protective information may be obtained from your cooperative Agricultural Extension Service.

### PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame

NOTICE: Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.

DEXOL INDUSTRIES Torrance, CA 90501

ł

្

EPA Reg. No. 192-142 EPA Est. 192-CA-1

BEST AVAILABLE COPY