

Reg # 192-142

PM-19

199

FRONT PANEL - CONTAINER LABEL

ACCEPTED
DEC 14 1993
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 192-142

DEXOL

DEXA-KLOR

INSECT SPRAY

Kills Ant, Fleas, Roaches, Chiggers, Ticks, Fire Ants, Grubs, Spiders, Mosquitoes, Mole Crickets, Grasshoppers, Crickets and More

Water-Based Formulation

1 Pint Treats Up to 2,666 Sq.Ft. of Lawn
OR
1 Quart Treats Up to 5,332 Sq.Ft. of Lawn

Lasts Up To 8 Weeks

Active Ingredient:

*Chlorpyrifos: O,O-diethyl O-(3,5,6-trichloro-2-
-2-pyridyl) phosphorothioate 6.7%


Inert Ingredients: 93.3%
Total: 100.0%

Keep Out of Reach of Children

C A U T I O N

See back panel for additional precautionary statement

*DURSBAN and the protection logo are trademarks of DowElance(Logo)



NET _____ FL.OZ. (_____ ML)

[CONTAINER LABEL - BACK PANEL]

SEE ATTACHED PAMPHLET FOR COMPLETE DIRECTIONS FOR USE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feedstuffs. Do not permit children or pets onto treated areas until the spray has dried completely. Do not contaminate ornamental fish ponds. Food utensils such as teaspoons or measuring cups should not be used for food purposes after use with pesticides.

STATEMENT OF PRACTICAL TREATMENT


If Swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious person. **If Inhaled:** Remove victim to fresh air. Apply artificial respiration if indicated and get medical attention. **If On Skin:** Remove contaminated clothing and wash affected areas with plenty of soap and water. **If In Eyes:** Flush eyes with plenty of water. Get medical attention if irritation persists. **Note to Physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection is the preferable antidote. Oximes, such as 2-PAM may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Protective information may be obtained from you Cooperative Agricultural Extension Service.

DEXOL INDUSTRIES
Torrance, CA 90501

EPA Reg. No. 192-142
EPA Est. No. 192-CA-1



[PAMPHLET LABEL]

DEXOL

DEXA-KLOR

INSECT SPRAY

Kills Ant, Fleas, Roaches, Chiggers, Ticks,
Fire Ants, Grubs, Spiders, Mosquitoes, Mole Crickets,
Grasshoppers, Crickets and More

Water-Based Formulation

1 Pint Treats Up to 2,666 Sq.FT. of Lawn
OR
1 Quart Treats Up to 5,332 Sq.Ft. of Lawn

Lasts Up To 8 Weeks

Active Ingredient:

*Chlorpyrifos: O,O-diethyl O-(3,5,6-trichloro-2-
-2-pyridyl) phosphorothioate 6.7%


Inert Ingredients: 93.3%
Total: 100.0%

Keep Out of Reach of Children

C A U T I O N

See back panel for additional precautionary statement

*DURSBAN and the protection logo are
trademarks of DowElnce(Logo)



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food and feedstuffs. Do not permit children or pets onto treated areas until the spray has dried completely. Do not contaminate ornamental fish ponds. Food utensils such as teaspoons or measuring cups should not be used for food purposes after use with pesticides.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious person. **If Inhaled:** Remove victim to fresh air. Apply artificial respiration if indicated and get medical attention. **If On Skin:** Remove contaminated clothing and wash affected areas with plenty of soap and water. **If In Eyes:** Flush eyes with plenty of water. Get medical attention if irritation persists. **Note to Physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection is the preferable antidote. Oximes, such as 2-PAM may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Protective information may be obtained from you Cooperative Agricultural Extension Service.

[THE FOLLOWING DIRECTIONS FOR USE WILL BE USED WITH PRODUCT PACKAGED "WITHOUT" A HOSE-END SPRAYER ATTACHMENT]

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is for use only in and around the home such as on lawns, outside the home and inside the home as directed. Do not apply this product to sites or areas not listed or prohibited on this label.

DO NOT APPLY THIS PRODUCT TO VEGETABLE GARDENS OR OTHER PLANTS INTENDED FOR FOOD OR FEED PURPOSES.

LAWN PEST CONTROL: For control of Ants, Armyworms, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Mole Crickets, Cutworms, Dichondra Flea Beetles, Earwigs, Fleas, Grasshoppers and Sod Webworms - Apply 3 fl.oz. per 15 gallons of water [1 fl.oz. or 2 tablespoons per 5 gallons water] to cover 500 sq.ft. of area. 1 pint of product will cover 2,666 sq.ft. of area.

[PAMPHLET LABEL - PAGE 3]

Spray when pests first appear. Repeat treatments as necessary. Thoroughly water treated area immediately except as noted. For best results, the lawn should be moist at time of treatment. One application will usually control Chinch Bugs for up to 8 weeks. For Sod Webworms, delay watering or mowing the treated area for 12 - 24 hours after treatment. For Mole Crickets, apply when early nymphs are active. Efficacy may be enhanced by applying late in the afternoon or early evening and watering the turf prior to treatment to move the mole crickets close to the soil surface. For Turfgrass (Hyperodis) Weevil, apply 6 fl.oz. per 15 gallons water [2 fl.oz. or 4 tablespoons per 5 gallons water] to cover 500 sq.ft. 1 pint of product will cover 1,333 sq.ft of area. Spray in mid-April and again in mid-May and follow recommended use directions. For White Grubs (European Chafer and Japanese Beetle larvae only), apply 6 - 12 fl. oz. per 30 gallons of water [1 - 2 fl.oz. or 2 - 4 tablespoons per 5 gallons water] to cover 500 sq.ft. 1 pint of product will cover 666 to 1,333 sq.ft. of area. Use the lower rate when infestations are light and the higher rate for severe infestations. Apply when grubs are young and actively feeding near the soil surface, usually late July and August. Immediately after spraying, water treated area with 1/2 to 2" of water. Consult your local Agricultural Extension Service Specialist for the proper time to treat in your area for Hyperodis Weevil and White Grubs. Damaged lawns and ornamental turf should be fertilized to stimulate re-growth and recovery from insect injury.

PESTS INDOORS: For spot treatment of Ants, Cockroaches, Clover Mites, Crickets, Firebrats, Silverfish and Spiders: Apply as a coarse spray or with a paint brush to corners, cracks in walls, along baseboards, beneath sinks, around stoves and plumbing. Use a 0.25% spray solution to control light infestations or a 0.50% spray solution to quickly reduce heavy infestations. To make a 0.25% spray solution, mix 5 1/2 fl.oz. in 1 gallon of water or 1 pint in 3 gallons of water. To make a 0.50% spray solution, mix 11 fl.oz. in 1 gallon of water or 1 quart in 3 gallons of water. For Ants, apply to ant trails, around doors and windows where entrance may occur. ALWAYS SPOT TEST AREAS TO BE TREATED FOR COLOR FASTNESS AND TO PREVENT STAINING BEFORE MAKING A COMPLETE APPLICATION. DO NOT APPLY THIS PRODUCT TO CONDUITS, MOTOR HOUSINGS, JUNCTION AND SWITCH BOXES OR OTHER ELECTRICAL EQUIPMENT BECAUSE OF POSSIBLE SHOCK HAZARDS.

OUTDOORS TO HELP PREVENT INFESTATION OF HOMES: To control Ants, Brown Dog Ticks, Chiggers, Cockroaches, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Flies, Grasshoppers, Millipedes, Mosquitoes, Sowbugs and Spiders - Apply 3 fl.oz. in 15 gallons of water [1 fl.oz. or 2 tablespoons in 5 gallons of water] as a spray or with a paint brush to outside surfaces of house, porches, screens, window frames, eaves, patios and garages. Repeat as needed. A perimeter spray may also be made by applying 2 fl.oz. in 12 gallons of water [1/2 fl.oz. or 1 tablespoon in 3 gallons of water] to a band of soil 6 to 10 feet wide around and adjacent to the home and 2 to 3 feet high on the foundation. 1 pint will cover 8,000 sq.ft. DO NOT ALLOW SPRAY TO CONTACT AZALEAS, CAMELLIAS, POINSETTIAS, ROSE BUSHES OR VARIEGATED IVY AS PLANT INJURY MAY BE OCCUR.

IMPORTED FIRE ANTS: Dilute 2 fl.oz. in 1 gallon of water. Apply 1 gallon of solution as a gentle rain to each mound using a sprinkle can. Thoroughly wet mound and surrounding area to a 4 foot diameter (12 1/2 sq.ft.). For best

[PAMPHLET LABEL - PAGE 4]

results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity and treat all colonies which may not as yet have constructed a mound.

PESTS OF ORNAMENTALS(Flowers, Shrubs, Vines, Shade or Flowering Trees and Evergreens) - NOTE: DO NOT USE THIS PRODUCT ON AZALEAS, CAMELLIAS, POINSETTIAS, ROSE BUSHES OR VARIEGATED IVY BECAUSE OF POSSIBLE INJURY TO THESE PLANTS. To control Aphid, Bagworm, Eastern Tent Caterpillar, Fall Webworm, Grasshopper, Orangestriped Oakworm, Yellownecked Caterpillar, Mites, Spittlebug and Whiteflies: Mix 6 fl.oz. in 9 gallons of water (4 teaspoons per gallon). Spray upper and lower leaf surfaces and infested limbs and trunk. Avoid spraying to the point of excessive run-off. Treat when pests appear and repeat at 7 to 10 day intervals if needed. **To control Cutworm, Leafhopper, Mealybug, Mimosa Webworm, Redhumped Caterpillar and Thrips (exposed):** Mix 12 fl.oz. in 9 gallons of water (8 teaspoons per gallon) and apply as directed above. **For control of Ash and Lilac Borers, Oak Pine Needle, Tea and White Peach Scale Crawlers:** Mix 12 fl. oz. in 12 gallons of water (1 tablespoon per gallon) and apply as directed above.

Note: 1 fl.oz. = 2 tablespoons
1 tablespoon = 6 teaspoons

STORAGE & DISPOSAL

Storage: Store in a cool, dry place. Avoid storing at temperatures below 32°F. Keep in a locked storage area away from children and pets. **Disposal:** Securely wrap original container in several layers of newspaper and discard in trash.


NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS AND PRECAUTIONS.

[THE FOLLOWING DIRECTIONS FOR USE WILL BE USED WITH PRODUCT PACKAGED "WITH" A HOSE-END SPRAYER ATTACHMENT]


DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is for use only in and around the home such as on lawns, outside the home and inside the home as directed. Do not apply this product to sites or areas not listed or prohibited on this label.

DO NOT APPLY THIS PRODUCT TO VEGETABLE GARDENS OR OTHER PLANTS INTENDED FOR FOOD OR FEED PURPOSES.


HOSE-END SPRAYER ATTACHMENT INSTRUCTIONS: 1) Remove cap and foil liner from bottle. 2) Place siphonage tube into bottle. 3) Turn sprayer off by turning on/off valve 1/4 turn. Securely tighten sprayer attachment to bottle. 4) Securely screw hose into hose-end sprayer attachment.




[PAMPHLET LABEL - PAGE 5]



SPRAYER INSTRUCTIONS: The hose-end sprayer attachment has been pre-calibrated to deliver 2 teaspoons per 1 gallon of water. 1) Turn on water. TO START FLOW OF PRODUCT, turn sprayer attachment on. Product will be automatically diluted at proper ratio. Hold about 3 feet above area to be treated moving at a steady pace to ensure consistent coverage. When applying, hold container by placing sprayer attachment in palm of hand and securely grasping container handle with fingers. 2) TO STOP FLOW OF PRODUCT, turn sprayer attachment off. NOTE: To ensure adequate coverage, use moderate to high water pressure when applying.



LAWN PEST CONTROL: For control of *Ants, Armyworms, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Mole Crickets, Cutworms, Dichondra Flea Beetles, Earwigs, Fleas, Grasshoppers and Sod Webworms* - Apply at a rate of 1 pint to cover 2,666 sq.ft. of area. Spray when pests first appear. Repeat treatments as necessary. Thoroughly water treated area immediately except as noted. For best results, the lawn should be moist at time of treatment. One



[PAMPHLET LABEL - PAGE 6]

application will usually control Chinch Bugs for up to 8 weeks. For Sod Webworms, delay watering or mowing the treated area for 12 - 24 hours after treatment. For Mole Crickets, apply when early nymphs are active. Efficacy may be enhanced by applying late in the afternoon or early evening and watering the turf prior to treatment to move the mole crickets close to the soil surface. For Turfgrass (Hyperodis) Weevil, apply at a rate of 1 pint to cover 1,333 sq.ft of area. Spray in mid-April and again in mid-May and follow recommended use directions. For White Grubs (European Chafer and Japanese Beetle larvae only), apply at a rate of 1 pint to cover 666 to 1,333 sq.ft. of area. Use the lower rate when infestations are light and the higher rate for severe infestations. Apply when grubs are young and actively feeding near the soil surface, usually late July and August. Immediately after spraying, water treated area with 1/2 to 2" of water. Consult your local Agricultural Extension Service Specialist for the proper time to treat in your area for Hyperodis Weevil and White Grubs. Damaged lawns and ornamental turf should be fertilized to stimulate re-growth and recovery from insect injury.

OUTDOORS TO HELP PREVENT INFESTATION OF HOMES: To control Ants, Brown Dog Ticks, Chiggers, Cockroaches, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Flies, Grasshoppers, Millipedes, Mosquitoes, Sowbugs and Spiders - Apply to wet outside surfaces of house, porches, screens, window frames, eaves, patios and garages. Repeat as needed. A perimeter spray may also be made by applying to a band of soil 6 to 10 feet wide around and adjacent to the home and 2 to 3 feet high on the foundation. Apply at a rate of 1 pint to cover 8,000 sq.ft. DO NOT ALLOW SPRAY TO CONTACT AZALEAS, CAMELLIAS, POINSETTIAS, ROSE BUSHES OR VARIEGATED IVY AS INJURY MAY BE OCCUR.

Dexol Dexa-Klor Insect Spray may also be used on the sites listed below. DO NOT use the nose-end sprayer attachment provided with this product. Use a commercially available adjustable hose-end or tank type sprayer unless otherwise directed below.

PESTS INDOORS: For spot treatment of Ants, Cockroaches, Clover Mites, Crickets, Firebrats, Silverfish and Spiders: Apply as a coarse spray or with a paint brush to corners, cracks in walls, along baseboards, beneath sinks, around stoves and plumbing. Use a 0.25% spray solution to control light infestations or a 0.50% spray solution to quickly reduce heavy infestations. To make a 0.25% spray solution, mix 5 1/2 fl.oz. in 1 gallon of water or 1 pint in 3 gallons of water. To make a 0.50% spray solution, mix 11 fl.oz. in 1 gallon of water or 1 quart in 3 gallons of water. For Ants, apply to ant trails, around doors and windows where entrance may occur. ALWAYS SPOT TEST AREAS TO BE TREATED FOR COLOR FASTNESS AND TO PREVENT STAINING BEFORE MAKING A COMPLETE APPLICATION. DO NOT APPLY THIS PRODUCT TO CONDUITS, MOTOR HOUSINGS, JUNCTION AND SWITCH BOXES OR OTHER ELECTRICAL EQUIPMENT BECAUSE OF POSSIBLE SHOCK HAZARDS.

IMPORTED FIRE ANTS: Dilute 2 fl.oz. in 1 gallon of water. Apply 1 gallon of solution as a gentle rain to each mound using a sprinkle can. Thoroughly wet mound and surrounding area to a 4 foot diameter (12 1/2 sq.ft.). For best results, apply in cool weather, 65-80°F, or in early morning or late evening.

999

[PAMPHLET LABEL - PAGE 7]

hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity and treat all colonies which may not as yet have constructed a mound.

PESTS OF ORNAMENTALS(Flowers, Shrubs, Vines, Shade or Flowering Trees and Evergreens) - NOTE: DO NOT USE THIS PRODUCT ON AZALEAS, CAMELLIAS, POINSETTIAS, ROSE BUSHES OR VARIEGATED IVY BECAUSE OF POSSIBLE INJURY TO THESE PLANTS. To control Aphid, Bagworm, Eastern Tent Caterpillar, Fall Webworm, Grasshopper, Orangestriped Oakworm, Yellownecked Caterpillar, Mites, Spittlebug and Whiteflies: Mix 6 fl.oz. in 9 gallons of water (4 teaspoons per gallon). Spray upper and lower leaf surfaces and infested limbs and trunk. Avoid spraying to the point of excessive run-off. Treat when pests appear and repeat at 7 to 10 day intervals if needed. **To control Cutworm, Leafhopper, Mealybug, Mimosa Webworm, Redhumped Caterpillar and Thrips (exposed):** Mix 12 fl.oz. in 9 gallons of water (8 teaspoons per gallon) and apply as directed above. **For control of Ash and Lilac Borers, Oak Pine Needle, Tea and White Peach Scale Crawlers:** Mix 12 fl. oz. in 12 gallons of water (1 tablespoon per gallon) and apply as directed above.
Note: 1 fl.oz. = 2 tablespoons
1 tablespoon = 6 teaspoons

STORAGE & DISPOSAL

Storage: Store in a cool, dry place. Avoid storing at temperatures below 32°F. Keep in a locked storage area away from children and pets. **Disposal:** Securely wrap original container and hose-end sprayer attachment in several layers of newspaper and discard in trash.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS AND PRECAUTIONS.

DEXOL INDUSTRIES
Torrance, CA 90501

EPA Reg. No. 192-142
EPA Est. No. 192-CA-1

