

DEC 18 1996

Mr. John Prues
Dexol Industries
1450 W. 228th Street
Torrance, CA 90501-5089

Dear Mr. Prues:

Subject: Eager Beaver Dormant Oil & Lime Sulphur Spray
EPA Reg. No. 192-201
Your Submission of November 6, 1996

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) for the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable provided that you:

1. Submit/cite all data required for registration and reregistration of your product under section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data.


2. Make the following labeling change: Change "Berries" to "Cane Berries" or indicate specific types of berries intended. The broad term "berries" would include strawberries, blueberries, and other types of plants but the pests indicated are those associated with raspberries, blackberries and other cane berries.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is included for your records.

Sincerely yours,



Connie B. Welch
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure
7505C:C.Grable:cg:12/6/96

2074

[Front Panel]

EAGER BEAVER BRAND
DORMANT
OIL AND LIME SULPHUR
SPRAY

For Dormant and Growing Season Disease Control

Controls: Red Spider Mites , Scale Insects
MealyBugs - Peach Leaf Curl

96 NOV 13 P1:43

RECD L/A/G/P/DPD1

ACTIVE INGREDIENTS:	BY WT.
Calcium Polysulfide.....	4.6%
Petroleum Oil.....	58.0%
INERT INGREDIENTS:	37.4%
TOTAL	100.0%

Unulfonated residue 92% minimum. Petroleum Oil Classification - Medium

KEEP OUT OF REACH OF CHILDREN

DANGER

ACCEPTED
with COMMENTS
In EPA Letter Dated

See Side Panel for Additional Precautionary Statements

NET CONTENTS

1 QUART (0.946 LITERS)

DEC 18 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

192-201

3044

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Corrosive to eyes. Concentrate causes eye damage and skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of mild, egg white, or gelatin solution or if these are not available, large quantities of water. **DO NOT INDUCE VOMITING.** Do not give anything by mouth to an unconscious person. Get medical attention. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention. **IF IN EYES:** Flush eyes with a gentle stream of water for 15 minutes. Get medical attention. **IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix with fertilizers, acid cleaning compounds or other acidic household chemicals. May give off highly toxic and extremely flammable hydrogen sulfide gas if mixed with acids.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DORMANT SPRAY is an excellent early spring "Clean-up" and dormant spray for CITRUS and many other FRUIT and SHADE TREES including, EVERGREEN AND DECIDUOUS SHRUBS and GARDEN PLANTS such as CAMELLIAS, AZALEAS, GARDENIAS, HIBISCUS, HOLLY, VIBURNUM, PRIVET and ROSES (after pruning). Dormant Spray controls Scale insects (Camellia, Rose, San Jose, Black, Red) in migratory state, MEALY BUGS, RED SPIDER MITES and PEACH LEAF CURL.

APPLICATION

Apply through hose end sprayer, tank pressure or trombone sprayer. Shake well before using. Partially fill sprayer with water and add the recommended amount of insecticide under agitation. Add balance of water. Apply using a coarse spray with considerable force. Thoroughly cover branches and trunks spraying to the point of runoff. Always prepare a fresh spray solution at time of use. Do not apply through any type of irrigation system.

Do not spray when temperature is expected to exceed 80° within 24 hours. Spray early in the morning or late in the afternoon to avoid burning. Make new dilution for each use. Do not use Dormant Season rates if leaves are present on plants as foliar injury may occur.

DORMANT SEASON SPRAYING

DO NOT APPLY IN SPRING AFTER BUDS BEGIN SHOWING NEW GROWTH. USE ONLY WHEN PLANTS ARE DECIDUOUS OR DORMANT. Shake well before using.

FRUIT TREES, ROSES, AND DECIDUOUS ORNAMENTALS & SHADE TREES (no leaves in winter): To control : San Jose Scale, Rose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, Overwintering Insect Eggs, Many overwintering Fungus Spores or Plant Diseases. Prune and destroy all old leaves and branches. Mix 8 tablespoons in one gallon of water (one pint in 4 gallons of water.)

4084

Specific Directions:

PEACHES: Peach leaf curl, Peach blight (Shot Hole Fungus)-For best results use a combination of late fall and mid-winter applications. LATE FALL-Spray in October or November after leaf drop, but before heavy rains occur. MID-WINTER-spray in January or February before buds begin to swell. If buds have begun to swell or open, it is too late to control Peach Leaf Curl Disease as infection has already occurred.

ROSES: Black Spot, Powdery Mildew, Rust-Prune and destroy all old leaves. Spray bushes in the spring before buds begin to swell.

PEARS: Blister Mites-Apply in the fall before the leaves start to turn color and again in the spring just as the buds begin to swell.

BERRIES: Leaf Spot, Cane Blight, Red Berry trouble (Recognized by Dwarf Berries)-Apply in the late fall followed by another treatment in spring as buds begin to swell.

GROWING SEASON SPRAYING

For Control of Scale Insects, Spider Mites, Mealybugs, and Whitefly larvae on Citrus (Oranges, Lemons, Grapefruit, Tangerines, Limes), Fruit & Nut Trees (Almonds, Apples, Plums, Pears, Apricots, Peaches, Cherries, Prunes, Nectarines, Olives), Shade Trees, Evergreens, Camellias, Azaleas, Hardy Shrubs. Use 3 teaspoons per gallon of water on tender new growth and 5 teaspoons per gallon of water on mature foliage. Apply every 1-2 weeks. NOTE: Early and mid season oranges may be injured from spray applied during summer and early fall.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place preferably in a locked storage area. Do not store diluted spray. **PRODUCT DISPOSAL:** Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse empty container.

NOTICE TO BUYER: Seller makes no warranty of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risk of using and handling.

Dexol Industries
1450 W. 228th St.
Torrance, CA. 90501

EPA REG. NO. 192-201
EPA EST. NO. 192-CA-1

Questions: (310) 326-8373

11/5/96