192 - 222



#### UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

apr 1 1 2011

4/11/2011

Ms. Diana Williams Value Garden Supply 9100 W. Bloomington Freeway, Suite 113 Bloomington, MN 55431

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Ms. Williams:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated March 18, 2011 for the following product(s):

AllPro Permethrin ICG .5% Reg. No. 192-222

The Registration Division (RD) has conducted a review of this request for applicability under PR Notice 2007-4 and finds that the label change(s) requested falls within the scope of PRN 2007-4. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please contact Linda A. DeLuise at 703-305-5428.

Sincerely,

Richard J. Gebken Product Manager Insecticide Branch Registration Division (7504P)

EPA	United States Environmental Protecti Washington, DC 2046	0 🛛 🖾 Other	ent	OPP Identifier Number	
	Applicat	ion for Pesticide -Section	nl		
1. Company/Product Number 192-222		2. EPA Product Manager R. Gebken	3. Proposed Classification ⊠ None □ Restricted		
4. Company/Product (Name) AllPro Permethrin ICC	9.5%	РМ# 10			
<ul> <li>5. Name and Address of Applicant (<i>Include ZIP Code</i>)</li> <li>Value Garden Supply</li> <li>P O Box 585</li> <li>St. Joseph, MO 64502</li> <li>Check if this is a new address</li> </ul>		6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.: Product Name:			
		Section I	<del>VPR i 1</del> 2011		
<ul> <li>Amendment - Explain Be</li> <li>Resubmission in respon</li> </ul>	low se to Agency letter dated	☐ Final printed labels in res _ ☐ "Me Too" Application	sponse to Agency	letter dated	
IX Notification - Explain be	ow	Other - explain below			
I. Material This Product Wil Child-Resistant Packaging ☐ Yes* ☐ No	alle i sheren en reest to et to te the second	Water Soluble Packaging       Pres       No	2. Type of Co	ntainer	
Certification must be submitted.	If "Yes," No. Per Unit Package Wt. Container	lf "Yes," No. Per Package Wt. Container	Glass Glass Paper Other (S	pecify)	
3. Location of Net Contents Information		4. Size(s) of Retail Container	<ul> <li>5. Location of Label Directions</li> <li>On Label</li> <li>On Labeling accompanying product</li> </ul>		
δ. Manner In Which Label Is A	Affixed To Product	lued			
		SectionIV			
. Contact Point (Complete ite	ms directly below for identification of i	ndividual to be contacted, if necessa	ry, to process this	s application.) o o	
<sup>Name</sup> Diana Williams		Title Reg. Affairs Mgr.			
	Certification ave made on this form and all attachm gly false or misleading statement may			6. Date Application	
2. Signature Kleana Willeam		3. Title Regulatory Affairs Manager		່ ຍ່ວວວ່ວ ມາລາວ ບ ຍາມກາ ບັນ ວ່ວຍວາມ	
I. Typed Name Diana Williams		5. Date		ι ι ι ι ι ι	
				Yellow - Applicant Co	



March 18, 2011

Mr. Richard Gebken Product Manager Team 10 Document Processing Desk (NOTIF) Insecticide Branch, Registration Division US EPA – OPP (7504P) 1200 Pennsylvania Ave., N.W. Washington, DC 20460

Subject: Notification of Revisions to Container Handling Label Instructions Per PR Notice 2007-4 EPA Reg. No. 192-222, AllPro Permethrin ICG .5% 3/8

9999 3 3393

ວ ວ ອອອອອອອ

Dear Mr. Gebken:

Enclosed for the subject registration are a Notification form and two copies of labeling revised to incorporate container handling and disposal instructions consistent with PR Notice 2007-4. The revisions are highlighted on one copy of the enclosed labeling.

If there are questions concerning the enclosed, please do not hesitate to contact me.

Sincerely,

Kleana William

Diana Williams Regulatory Affairs Manager On Behalf of Value Garden Supply

Enclosures

Cc: B. Russell, VGS

# AllPro<sup>®</sup> Permethrin ICG .5%

Kills insects in lawns

 $+\infty$ 

۰, <sup>3</sup>

- Lawn & Garden Insect Control
- Treat up to 5, 000 10,000 15,000 sq. ft.
- Controls home invading pests
- 10 pounds treats 5, 000 sq. ft.
- Controls surface and foliage feeding pests
- For grass and dichondra lawns
- Contains synthetic pyrethroid
- For outdoor use on home lawns
- Contains permethrin
- Ready to use
- Provides contact residual control for up to 4 weeks

Kills – Ants, Armyworms (including Fall, Yellow striped, etc.), Bear Tick, Bermuda grass Mites, Billbugs, (adult), Black Legged Tick, Boxelder Bugs, Brown Dog Tick, Centipedes, Chiggers, Chinch Bugs, Clover Mites, Cockroaches, Crickets, Cutworms, Darkling Ground Beetles, Deer Ticks, Digger wasps, Essex Skipper, European Crane Fly Larvae, Field Crickets, Fiery Skipper, Fire Ants, Flea Beetles, (in Dichondra), Grasshopper, Hyperodes (Turfgrass Weevil), Japanese Beetle (adult), Lawn Moth, Leaf Hoppers, Lone Star Tick, Scorpions, Silverfish & Firebrats, Sod Webworms, Sowbugs, Spiders (Including Black Widow), Spittlebugs, Springtails, Vegetable weevils (in Dichondra), Western Black Legged Tick (*Ixodes pacificus*)

NOTIFICATION

APR 1 1 2011

ACTIVE INGREDIENT:	
Permethrin*	0.5%
INERT INGREDIENTS	<u>99.5%</u>
TOTAL	

\*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2, 2-dichlororethenyl) 2, 2 dimethyl cyclopropanecarboxylate. cis/trans ratio: maximum 55% (+/-) cis and minimum 45% (+/-) trans.

## KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No.192-222			EPA Est. No. 446	3 3 3 J	
	Value Garden PO Box 58			2200 3000 3000 3000 3000	ວງວອງອງ 
	St. Joseph, MO	64501			193939 00000 00000
	Net Content:	lbs.		ن وودودر ر ر ر ر ر ر ر ر ر ر ر ر ر ر ر ر	22272
				- 1 -	

	FIRST AID
	(Pyrethroid)
IF IN EYES	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
IF ON SKIN OR CLOTHING	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED	<ul> <li>Call a poison control center or doctor for immediate treatment advise.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a position control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
	HOT LINE NUMBER (Pyrethroid)

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

### **ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish and other aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment wash waters.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its EPA labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

READ ENTIRE LABEL BEFORE USING, USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

AllPro Permethrin ICG .5% is recommended for control of lawn insects, fleas, ticks outdoors in residential areas. Repeated applications are made by broadcast treatment with a spreader to lawns, under shrubbery, and other ornamental and turf grass areas, band treatments are recommended around the house foundation and spot applications are made to ant mounds and wasp nests.

## **General Use Directions**

- 1. Determine the size of the area to be treated measure and multiply length times width.
- 2. Determine appropriate spreader setting using chart.
- 3. treat as directed Through coverage is important.
- 4. Water in with 1/4 to 1/2 inches of sprinkler irrigated water to release the insecticide from the granule where indicated.

- 5. Keep children and pets off treated areas until this material is washed into turf and grass is dry
- 6. Do not use food utensils such as teaspoons or measuring cups for food purposes after use with pesticides

#### **Overall Lawn Treatment for Grass and Dichondra Lawns**

To control: Ants, Armyworms (including Fall, Yellow striped, etc.), Bear Tick, Bermuda grass Mites, Billbugs, — (adult), Boxelder Bugs, Brown Dog Tick, Centipedes, Chiggers, Chinch Bugs, Clover Mites, Cockroaches, Crickets, Cutworms, Darkling Ground Beetles, Deer Ticks, Digger wasps, Essex Skipper, European Crane Fly Larvae, Field Crickets, Fiery Skipper, Fire Ants, Flea Beetles, (in Dichondra), Grasshopper, Hyperodes (Turfgrass Weevil), Japanese Beetle (adult), Lawn Moth, Leaf Hoppers, Lone Star Tick, Scorpions, Silverfish & Firebrats, Sod Webworms, Sowbugs, Spiders (Including. Black Widow), Spittlebugs, Springtails, Ticks, Vegetable weevils (in Dichondra).

Apply this product uniformly to home lawns, under shrubbery, and other ornamental recreational turf grass areas. Treat when insects first appear using granules at the rate of 2 pounds per 1, 000 sq. ft. using a fertilizer spreader. See spreader setting chart below. To help obtain uniform distribution, apply one-half required amount of product in one direction, then apply the balance at a right angle. Immediately after application, water the lawn lightly. Retreat after 7 days if insects are present or damage reoccurs.

European Crane fly larvae (Leatherjackets) – the best time to apply is in May or June when the insects are young.

Chinch Bugs – Start treatments when temperature reaches 80°F or above. A second application may be necessary in 6 weeks if insect activity continues.

#### Fleas and Tick Control (Outdoors)

In summer months, fleas and ticks frequently become established in lawns and pet resting areas surrounding the home. For best control of fleas and ticks (including but not limited to Bear Tick, Black-legged Tick, Brown Dog Tick, Deer Tick (*Ixodes dammini*), Lone Star Tick, Western Black-legged Tick, Wood Tick); a regular treatment program should be followed, Treat once a month when pests first appear in late Spring (May – June) and continue until late Summer (September – October). Fall temperatures below 50 °F decrease pest activity. Eliminating weeds around the home and mowing lawns frequently reduces the hiding places for these pests.

Note: Dogs, cats, their kennel or bedding area should also be treated with a registered product. In addition, if fleas and ticks are indoor on carpets or pet resting areas, use a registered product to treat these areas. It is very important that all three sites – outdoors, indoors and also the pet itself be treated with appropriate labeled products to achieve satisfactory control. If any of these areas are not treated it could serve as a source of reinfestation to the entire premise.

#### Sod Webworms (Crambus spp.)

Sod webworms are light brown caterpillars 1/4 to 3/4 inches long that hide in the grass during the day and feed on the grass blades at night. Adult Webworms, usually referenced to a Lawn Moths ( about 1 inch in length with long snout) emerge from the grass at dusk. They fly over the lawn laying eggs that later hatch into destructive caterpillars. Damage appears first as small brown spots which, if not treated, may rapidly develop into large brown areas of severely damaged or dead grass. If Lawn Moths (Sod Webworm adults) are noticed in large numbers at dusk, or at first sign of damage to lawn, treat entire lawn evenly.

#### Band (Perimeter) Treatment Around House Foundation

To help prevent infestation of building by Ants, Boxelder Bugs, Brown Dog ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Fleas, Scorpions, Sowbugs, Spiders (including Black Widow), Springtails (*Collembola*) and Ticks. Apply a 5-foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly after application. Retreat after 7 days if insects are present.

#### Spot Treatment

Ants Nesting Outdoors – This product maybe used outdoors for the control of ants, including Acrobat, Argentine, Big Headed, Cornfield, Field, Harvester, Leafcutter, Little Black, Pavement and Pyramid Ants. If possible, it is best to try to locate nest and treat this first. Nests will usually be found in the ground but may be concealed under stones, boards, slabs, or in stumps or logs. Product should be applied to thoroughly cover the nest (1 to 2 teaspoons) and sprinkled gently with water. In addition to treating ant nests or mounds, entry points of buildings (used by ants) and the soil surface along the perimeter of the building should be treated,

Digger Wasps: Apply 1 to 2 teaspoons of granules in and around each individual ground nest opening. Make application at dusk when the wasps are not active.

Imported Fire Ants: Apply one cup over, and 2 feet around each fire ant mound: Immediately after application, water the granules into the soil gently with a light sprinkling to avoid disturbing the colony. Use at least one gallon of water per mound, but no more than 2 gallons. Use of high pressure watering equipment may disturb the colony and cause migration of the ants, thus reducing the effectiveness of the treatment. Apply in cool weather (65 to 80 °F) or in early morning or late evening for best results. Treat new mounds as they appear.

Spreader Make and Model	2 lbs/1000 sq. ft.	3 lbs/1000 sq. ft.
Central Imperial No. 1662-G	6½	71/4
Central Suburbia Lawn Spreader No. 1622-0	6½	71/4
Cyclone Model B-1 Rotary	41/2	5
Greenfield Model #105	61⁄2	71/4
Ortho Drop Spreader Model 3000	41/2	51/2
Ortho Broadcast Spreader Model 4000	41/2	7
JC Penney Model 6005	71/4	8
JC Penny Rotary Spreader	5	7
Scotts Model 35	4	5
Scotts Model PF 3	F	Н
Sears Craftsmen Model 671.19198	6	7
VW&RMasterlineR75FB	F	Н

#### SPREADER SETTINGS

Note: Spreader settings are approximate and variation can occur between identical models. User should calibrate spreader by accurately weighing a few pounds into your spreader and applying to a measured area. Then adjust spreader setting if necessary to assure correct rate of application.

## STORAGE AND DISPOSAL

**STORAGE**: Store product in its original container (keep closed), in a locked, cool and dry area out of reach of children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Non-refillable container. Do not reuse or refill container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste authority for disposal instructions. Never place unused product down any indoor or outdoor drain.

DISCLAMIER: Value Garden Supply warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Value Garden Supply makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent Supply shall not be liable for any consequential, special, or indirect damages resulting from the use, handling or application of this product.

• •

3 18 11 S&D Notif.

..

 $\sim 10^{10}$ 

8/8