48 /3

SHELL CHEMICAL, COMPANY
A Division of Shell Oil Company
Agricultural Chemicals Division

New York 20, New York
April 13, 1964

SPECIMEN LABEL

(YOUR DRAND NAME) Technical BIDRIN® Insecticide (Contains 9.0 lb BIDRIN per Gallon)

WEIGHT
90.0
LO.O 00.0%

GENERAL DIRECTIONS

Use this product for control of Mimosa Webworm on honey locust trees only at recommended dosages. To be used by professional arborists and nurserymen who have been specifically trained in the application of BIDRIN $^{\odot}$ Insecticide as a tree root implant.

USE AND DIRECTIONS FOR APPLICATION

For control of Mimosa Webworm on honey locusts by root implantation of BIDRIN Insecticide depending on the size and shape of a tree's crown, several large roots should be selected and uncovered for implanting BIDRIN Insecticide. Holes should be drilled at least one inch deep in the roots below the soil line with a 1/4 to 3/8 inch bit. Care should be taken to insure that the holes do not go completely through the roots. The recommended amount of BIDRIN, 1 to 2 ml for each inch of tree diameter for trees 6 inches or less in diameter and 2.5 to 3 ml per inch of diameter for trees larger than 6 inches in diameter, should be equally apportioned into the holes with a suitable instrument such as an automatic pipette. Soil removed when the roots were exposed for drilling should be replaced when the treatment is completed. This application technique is for use by professional arborists and nurserymen who have been specifically trained in root implantation of BIDRIN Insecticide.

PRECAUTIONS IN USING

Keep out of the reach of children and domestic animals. Do not contaminate food or feed products. Keep away from heat and open flame. To prevent re-use, empty containers should be decontaminated and destroyed. Decontaminate containers by washing thoroughly with strong solution of alkali and detergent and rinse with water.

492/3

PRECAUTIONS IN USING (Cont'd.)

DANGER! POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO BIDRIN POISONING.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors. Wear clean natural rubber gloves, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. During commercial or prolonged exposure in spray-mixing and loading operations, wear a mask or respirator of a type adequate against BIDRIN vapors and gases. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least ten minutes and get medical attention. In case of accidental contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with concentrated solution of alkali and detergent and rinse with water.

NOTE: Caution! Avoid skin and eye contamination when handling strong caustic solutions.

POISON (deg)

ANTODOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR BIDRIN POISONING. Consult your physician about obtaining an adequate supply of 1/100-grain atropine tablets for emergency use.

CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING

INTERNAL: If the material has been swallowed, induce vomiting immediately.

This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water).

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the BIDRIN, remove all contaminated clothing, and wash skin with scap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes. IF WARNING SYMPTOMS APPEAR: ADMINISTER two 1/100-grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR (See WARNING SYMPTOMS below). Keep-patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against para-sympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams, until complete a tropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

NOTICE OF WARRANTY

(Your Company Name) warrants (its liability being limited to the purchase price of the product named on this label) that this product consists of the ingredients specified but makes no other warranty or representation of any kind, expressed or implied, including any warranty or representation regarding the effect or result of this product's use, whether or not the product is used in accordance with directions, and shall have no responsibility whatsoever for injury to persons or loss or damage to property, resulting from the handling, storage, or use of this product. The user or buyer shall be deemed to have accepted these conditions, which may be varied only by agreement in writing, signed by duly authorized representatives of the above-named company.



MAY **26** 1964

UNDER THE FEDERAL INSECTIONE FOR GREIDE AND RODENT CHOIC ACT FOR ECONOMIC POISON REGISTERED UNDER NO.